

Finite groups of bounded rank with an almost regular automorphism. *

A. Jaikin-Zapirain

Departamento de Matemáticas-Matematika Saila

Facultad de Ciencias-Zientzi Fakultatea

Universidad del País Vasco-Euskal Herriko Unibertsitatea

Apdo. 644, 48080-Bilbao, Spain

Abstract

In this paper we prove that any finite group of rank r , with an automorphism whose centralizer has m points, has a characteristic soluble subgroup of (m, r) -bounded index and r -bounded derived length. This result gives a positive answer to a problem raised by E. I. Khukhro and A. Shalev (see also Problem 13.56 from the “Kourovka Notebook” [5]).

1 Introduction.

There has been certain interest on the study of finite groups with an automorphism of some fixed type over the last years. The classical restrictions on an automorphism consist in fixing the order of the automorphism and the order of the centralizer. We refer the interested reader to [4] for background on this subject.

In [8] A. Shalev began a new approach in this area. The **rank** of a group G (denoted by $\text{rk } G$) is the minimal integer r such that every subgroup of G is r -generated, and the **rank** of a \mathbb{Z}_p -Lie ring L is the minimal number of generators of L as \mathbb{Z}_p -module. In his work Shalev proved that if a finite group of rank r has an automorphism whose centralizer has m elements, then it has a soluble subgroup of (m, r) -bounded index. (In this paper we say that a certain invariant is (a, b, \dots) -bounded if it is bounded above by some function of (a, b, \dots) .) If, in addition, the orders of the automorphism and the group are coprime then the derived length of the subgroup can also be bounded by some function of (m, r) . The key to Shalev’s proof is the reduction of the problem to the analog for uniform \mathbb{Z}_p -Lie rings of bounded rank, which is valid without any hypotheses on the orders. In [3] E.I. Khukhro, assuming also that the order of group and the

*This work has been supported by a grant of the Government of the Basque Country and by the DGICYT Grant PB97-0604.

order of automorphism are coprime, improved Shalev's result, showing that the derived length of some soluble subgroup of (m, r) -bounded index is r -bounded.

In this paper we consider the general case and we prove the following result.

Theorem 1.1. *Let G be a finite group of rank r admitting an automorphism with m fixed points. Then G has a characteristic soluble subgroup H , whose index is (m, r) -bounded and whose derived length is r -bounded.*

We will call a Lie ring L a **uniform** Lie ring if it is a finitely generated free $\mathbb{Z}/p^i\mathbb{Z}$ -module for some i and some prime p . In order to prove Theorem 1.1 we use the reduction from [9] to uniform Lie rings. In Section 2 we consider the Lie ring case and in Section 3 we give the proof of Theorem 1.1 in the case of p -groups. In Section 4 we finish the proof of Theorem 1.1, following the lines of [8, 3] and also put some conjectures. The notation is standard. The derived series of a group is denoted by $\{G^{(k)}\}$ and a similar notation will be used for Lie rings. We will denote by G^n the subgroup of a group G generated by all n th powers of elements in G . $[r]$ denotes the upper integral part of a real number r .

2 Lie ring case

Let \mathbb{Z}_p be the ring of p -adic integers and \mathbb{Q}_p its field of quotients. Set $S = \mathbb{Z}_p[x, x^{-1}]$. For every finitely generated S -module M such that $M/(x-1)M$ is finite, define the x -**rank** of M , $\text{rx } M = \log_p(|M : (x-1)M|)$. Note that if M is finite then $|C_M(x)| = p^{\text{rx } M}$. Throughout this paper we shall call Lie \mathbb{Z}_p -(sub)algebras simply **Lie (sub)rings** for brevity and then Lie automorphism will mean Lie \mathbb{Z}_p -algebra automorphism. The following theorem is the main result of this section.

Theorem 2.1. *There is a function $f = f(p^m, r)$ such that if G is a Lie ring of rank r and also a finitely generated S -module of x -rank m with x operating on it as a Lie automorphism, then G has a soluble subring H of index less than f and derived length at most 1 if $r = 1$ and $2^r - 2$ if $r > 1$. Moreover, if G is a uniform Lie ring then the derived length of H is at most 2^{r-1} .*

In order to prove this theorem we need some preliminary work. In this paper all the tensor and exterior products are taken over \mathbb{Z}_p . Let L be an S -module. Then we can define in $L \wedge L$ an S -module structure by setting

$$x(a \wedge b) = (xa) \wedge (xb), \quad x^{-1}(a \wedge b) = (x^{-1}a) \wedge (x^{-1}b) \quad \text{where } a \wedge b \in L \wedge L.$$

We define the category \mathbb{K} , whose objects are the triples $(L, +, \cdot)$ where

- (i) $(L, +)$ is an S -module.
- (ii) \cdot is a \mathbb{Z}_p -bilinear and alternating composition in L (i.e. \cdot defines an element of $\text{Hom}_{\mathbb{Z}_p}(L \wedge L, L)$).

The morphisms of \mathbb{K} are the S -homomorphisms preserving the multiplication. In the sequel the elements of \mathbb{K} will be called **rings** and we shall write (L, α) instead of L when we want to emphasize that the multiplication \cdot in a ring L is given by $\alpha \in \text{Hom}_{\mathbb{Z}_p}(L \wedge L, L)$. For any $L_1, L_2 \in \mathbb{K}$ we shall write $L_1 < L_2$ if L_1 is a proper subring of L_2 , i.e. L_1 is a proper subset of L_2 and the inclusion of L_1 into L_2 is a morphism in \mathbb{K} . We shall use \cong for isomorphism of \mathbb{Z}_p -modules and $\cong_{\mathbb{K}}$ for isomorphism in the category \mathbb{K} .

If $L \in \mathbb{K}$ and $A, B \subseteq L$ let $A \cdot B$ be the S -submodule generated by $a \cdot b$, where $a \in A$ and $b \in B$. Let $\Gamma_1(L)$ be the S -submodule of L generated by the elements

$$a \cdot (b \cdot c) + c \cdot (a \cdot b) + b \cdot (c \cdot a), \quad x(a \cdot b) - (xa) \cdot (xb), \quad a, b, c \in L,$$

and, for $i > 1$, let $\Gamma_i(L) = \Gamma_{i-1}(L) \cdot L$. Set $\Gamma(L) = \sum_i \Gamma_i(L)$ (so $\Gamma(L)$ is the ideal of L generated by $\Gamma_1(L)$). It is clear that $\bar{L} = L/\Gamma(L)$ becomes a Lie ring if for $\bar{a}, \bar{b} \in \bar{L}$ we define its Lie bracket by $[\bar{a}, \bar{b}] = (a \cdot b) + \Gamma(L)$, and x acts on \bar{L} as a Lie automorphism. We call $L \in \mathbb{K}$ a **lattice** if $L \cong (\mathbb{Z}_p)^s$ for some s . The key to the proof of Theorem 2.1 is to prove it for the Lie rings $L/\Gamma(L)$ when L is lattice (see Theorem 2.4). Recall the following lemma proved in [2].

Lemma 2.2. *Let L be an S -module and finitely generated as \mathbb{Z}_p -module. If the x -rank of $L/p^{m+1}L$ is $\leq m$ then the x -rank of L is also at most m .*

Lemma 2.3. *Let A be an S -module of x -rank m , finitely generated as a \mathbb{Z}_p -module, and let B be an S -submodule of A . Then $\text{rx } B \leq m$.*

Proof. Remember that the case when B is of finite index in A was proved in [2, Lemma 3.6]. Let $\bar{B} = \{a \in A \mid p^k a \in B \text{ for some } k \geq 0\}$. Then, by [2, Lemma 3.6], $\text{rx } B \leq \text{rx } \bar{B}$ and $\text{rx}(\bar{B} + p^{m+1}A)/p^{m+1}A \leq \text{rx } A/p^{m+1}A \leq m$. It directly follows from the definition of \bar{B} that $\bar{B} \cap p^{m+1}A = p^{m+1}\bar{B}$. Hence $\text{rx } \bar{B}/p^{m+1}\bar{B} \leq m$, and, by Lemma 2.2, $\text{rx } \bar{B} \leq m$. \square

Theorem 2.4. *Let G be a Lie ring of rank r and also a finitely generated S -module of x -rank m with x operating on it as a Lie automorphism. Suppose that $\text{rk } p^m G = r$. Then there exists a lattice $L \in \mathbb{K}$ of rank r and x -rank m , such that G is an epimorphic image (as an element of \mathbb{K}) of $\bar{L} = L/\Gamma(L)$.*

Proof. Let $L = (\mathbb{Z}_p)^r$ and let β be a surjective morphism from L onto G . Using that L is a free \mathbb{Z}_p -module, we obtain that there exists $\phi \in \text{End}_{\mathbb{Z}_p}(L)$ such that $x\beta(l) = \beta(\phi(l))$ for every $l \in L$. Since x induces a bijection on G/pG , ϕ induces a bijection on L/pL and so ϕ is a bijection. Define on L a structure of S -module by way of $xl = \phi(l)$, $x^{-1}l = \phi^{-1}(l)$ for every $l \in L$. The condition $\text{rk } p^m G = r$ implies that $\text{Ker } \beta \subseteq p^{m+1}L$ and so $|L : ((x-1)L + p^{m+1}L)| \leq p^m$. According to Lemma 2.2, the x -rank of L is at most m .

Since $L \wedge L$ is \mathbb{Z}_p -free, there exists $\alpha \in \text{Hom}_{\mathbb{Z}_p}(L \wedge L, L)$ such that $\beta(\alpha(l \wedge m)) = [\beta(l), \beta(m)]$ for every $l, m \in L$. It is clear now that $\beta(\Gamma(L, \alpha)) = 0$, and so G is an epimorphic image of $\bar{L} = L/\Gamma(L, \alpha)$. \square

Thus, we see that in order to prove Theorem 2.1 there is no loss of generality if we take G to be a Lie ring of the type $\bar{L} = L/\Gamma(L)$ where L is a lattice.

Define $\mathbb{E} = \{L \in \mathbb{K} \mid L \cong (\mathbb{Z}_p)^r, \text{rx } L = m\}$. If $L \in \mathbb{E}$, the tensor product $\mathbb{L} = \mathbb{Q}_p \otimes L$ belongs to the category \mathbb{K} . We call \mathbb{L} **simple** if $\mathbb{L}^2 \neq 0$ and there is no proper $\mathbb{Q}_p[x, x^{-1}]$ -submodule A of \mathbb{L} such that $\mathbb{L} \cdot A \subseteq A$. Also a lattice M is called **maximal** if there are no lattice $N \in \mathbb{E}$ and injective morphism $\phi: M \rightarrow N$ in \mathbb{K} such that $\phi(M) < N$. The proof of the following result is the same as the one of [2, Corollary 4.6]. We include the main steps of it for the sake of completeness.

Lemma 2.5. *Let $L \in \mathbb{E}$ and let \mathbb{L} be simple. Then there exists a maximal lattice $M \in \mathbb{E}$, such that $L \leq M < \mathbb{L}$.*

Proof. We split the proof in a number of steps.

Step 1. Let N be a lattice and $L < N < \mathbb{L}$. Define $t(L) = \min\{t \mid p^t L \subseteq L^2\}$ and $k = \max\{l \mid L \subseteq p^l N\}$. Then $t(L)$ and k are finite and $k \leq t(L)$.

Since \mathbb{L} is simple, $\mathbb{L}^2 = \mathbb{L}$ and $\mathbb{Q}_p L^2 = \mathbb{L}^2 = \mathbb{L}$. Hence $|L : L^2|$ is finite and so $t(L)$ is also finite. Using that N/L is a torsion finitely generated \mathbb{Z}_p -module, we also obtain that k is finite. By the definitions of $t(L)$ and k , we have

$$p^{t(L)} L \subseteq L^2 \subseteq p^{2k} N^2 \subseteq p^{2k} N.$$

Therefore by the maximality of k it follows that $2k - t(L) \leq k$, whence $k \leq t(L)$.

Step 2. There is no proper ascending series of lattices $L < L_1 < L_2 \dots < \mathbb{L}$.

Otherwise, put $N = \cup_{i \geq 1} L_i$ and define $A = \{a \in N \mid p^{-k} a \in N \text{ for every } k \in \mathbb{N}\}$. Then A is a $\mathbb{Q}_p[x, x^{-1}]$ -submodule of \mathbb{L} . For any $l \in N$, $a \in A$ and $k \in \mathbb{N}$ we have

$$p^{-k}(l \cdot a) = l \cdot (p^{-k} a) \in N.$$

Hence $N \cdot A \subseteq A$. Since $\mathbb{L} = \mathbb{Q}_p N$, $\mathbb{L} \cdot A \subseteq A$ and either $A = 0$ or \mathbb{L} .

In the former case, by [2, Lemma 4.3], N is a finitely generated \mathbb{Z}_p -module and so, $|N : L|$ is finite, which is a contradiction.

If $A = \mathbb{L}$ fix a_1, \dots, a_m an \mathbb{Z}_p -system of generators of L . Since $p^{-t(L)-1} a_i \in N$, there exists $k \geq 1$ such that $p^{-t(L)-1} a_i \in L_k$ for all i . Hence $L \subseteq p^{(t(L)+1)} L_k$, which contradicts Step 1.

Step 3. There exists a maximal lattice M such that $L \leq M < \mathbb{L}$.

By the previous step, there exists a lattice M such that $L \leq M < \mathbb{L}$ and M is maximal with this property. We shall prove that M is a maximal lattice. Suppose by way of contradiction that there exist a lattice N and an injective morphism ϕ such that $\phi(M) < N$ and $M \cong N$. Since N and $\phi(M)$ have the same rank, $N/\phi(M)$ is finite and so for every $n \in N$ there exists $k \in \mathbb{N}$ such that $p^k n = \phi(m)$ for some $m \in M$. We define $\psi: N \rightarrow \mathbb{L}$ by means of $\psi(n) = p^{-k} m \in \mathbb{L}$. This map is well defined and it is a morphism in the category \mathbb{K} . Hence $L \leq M < \psi(N) < \mathbb{L}$, against the choice of M . \square

We shall need the following result on finite dimensional Lie algebras from [6].

Proposition 2.6. *Let \mathbb{F} be a field, L a finite dimensional \mathbb{F} -Lie algebra and ϕ an automorphism of L . If the centralizer $C_L(\phi)$ is trivial, then L is soluble of derived length at most $\dim_{\mathbb{F}} L$.*

Let $L \in \mathbb{E}$. Define $\gamma(L) = \sup\{k \mid \Gamma_1(L) \subseteq p^k L\}$. Put $L^{(0)} = L$ and for $i > 0$, $L^{(i)} = L^{(i-1)} \cdot L^{(i-1)}$.

Lemma 2.7. *There exists $a = a(p, m, r) \geq 0$ such that for every maximal lattice $L \in \mathbb{E}$ and for every $s \geq 0$, $\gamma(p^s L) \leq 2s + a$.*

Proof. Suppose that for every j there exists a maximal lattice $M \in \mathbb{E}$ such that $\gamma(M) > j$. Then the set of Lie rings $T_0 = \{M/p^{\gamma(M)}M \mid M \in \mathbb{E} \text{ is maximal}\}$ is infinite. Now, suppose that we construct a Lie ring $N_k \in \mathbb{K}$, such that $N_k \cong (C_{p^k})^r$ and the subset $T_k = \{N \in T_0 \mid N/p^k N \cong_{\mathbb{K}} N_k\}$ of T_0 is infinite. Then there is a Lie ring $N_{k+1} \in \mathbb{K}$ such that $N_{k+1} \cong (C_{p^{k+1}})^r$ and the subset $T_{k+1} = \{N \in T_k \mid N/p^{k+1} N \cong_{\mathbb{K}} N_{k+1}\}$ of T_0 is infinite. Following these constructions we obtain a series $\{N_i\}$ of Lie rings, such that $N_i \cong_{\mathbb{K}} N_{i+1}/p^i N_{i+1}$. Let L be an inverse limit of $\{N_i\}$. Then L is a Lie ring and is also an S -module, where x acts as a Lie automorphism. Since $\text{rx } L/p^{m+1}L = \text{rx } N_{m+1} = \text{rx } N/p^{m+1}N$ for any $N \in T_{m+1}$, by Lemma 2.2, $\text{rx } L = m$ and so x acts without fixed points on L . By Proposition 2.6, $\mathbb{Q}_p \otimes L$ is soluble, whence there is an abelian ideal $A \neq 0$ of $\mathbb{Q}_p \otimes L$ which is also a $\mathbb{Q}_p[x, x^{-1}]$ -submodule. Put $B = A \cap L$. It is clear that B is an abelian ideal of L , an S -submodule of L and that $p^{-1}B \not\subseteq L$. Take any maximal lattice M such that $M/p^{\gamma(L)}M$ belongs to T_2 . Hence $M/p^2M \cong_{\mathbb{K}} N_2$. Let C be the image of B in $N_2 \cong L/p^2L$ and let D be the preimage of C under the natural projection from M to $N_2 \cong M/p^2M$. By the construction, D is an S -submodule of M , $D \cdot D \subseteq p^2M$, $M \cdot D \subseteq D + p^2M$ and $p^{-1}D \not\subseteq M$. Put $P = M + p^{-1}D$. Consequently, $P \cdot P \subseteq P$ and $M < P$, a contradiction with the maximality of M .

Hence there exists $a = a(p, m, r)$ such that $\gamma(M) \leq a$ for every maximal lattice $M \in \mathbb{E}$. Since $p^{3s}\Gamma_1(M) \subseteq \Gamma_1(p^s M)$, we obtain that $\gamma(p^s M) \leq 2s + a$. \square

Now we complete the proof of Theorem 2.1.

Proof of Theorem 2.1. We shall prove Theorem 2.1 by induction on r . It is obvious that Theorem 2.1 is true for $r = 1$ and $r = 2$. Suppose now that $r > 2$.

Consider the case $m = 0$. We can suppose that G is finite because it is approximated by $G/p^s G$, $s \geq 1$. Let $n = n_1 p^k$ where n_1 and p are coprime and x^n acts trivially on G . Note that the condition $m = 0$ implies $C_G(x) = 0$. Now, $(G, +)$ is a finite p -group and $C_G(x) = C_{C_G(x^{p^k})}(x)$, therefore $C_G(x^{p^k}) = 0$. Accordingly, in this case Theorem 2.1 follows from [10, Proposition 6.8].

Now suppose that $m > 0$. Using Theorem 2.4, we may assume that $G \cong_{\mathbb{K}} L/\Gamma(L)$, where $L \cong \mathbb{Z}_p^r$ is a lattice of x -rank m . Moreover, if G is uniform then $G \cong_{\mathbb{K}} L/p^{\gamma(L)}L$. First, suppose that $\mathbb{L} = \mathbb{Q}_p \otimes L$ is simple. By Lemma 2.5, there exists a maximal lattice M such that $L \leq M < \mathbb{L}$. We define $s = \min\{k \mid$

$p^k M \subseteq L\}$. Hence the rank of $L/((p^{s-1}M \cap L) + \Gamma(L))$ is less than r and so, by the induction hypothesis, we obtain that there is a function $h = h(p, m, r)$ such that $(p^h L)^{(2^{r-1}-2)} \subseteq p^s M + \Gamma(L)$.

If $\Gamma_1(L) \subseteq p^{\gamma(p^s M)+s+1}L$, then

$$\Gamma_1(p^s M) \subseteq \Gamma_1(L) \subseteq p^{\gamma(p^s M)+s+1}L \subseteq p^{\gamma(p^s M)+1}p^s M.$$

It is impossible, whence $\gamma(L) \leq \gamma(p^s M) + s$. Applying Lemma 2.7, we obtain $\gamma(L) \leq 3s + a$. In particular, $p^a(p^s M)^{(2)} \subseteq p^{\gamma(L)}L$ and, consequently, $(p^g L)^{(2^{r-1})} \subseteq p^{\gamma(L)}L$ for some $g = g(p, m, r)$. If G is uniform then we are done. The general case follows from the inductive hypothesis.

Suppose now that \mathbb{L} is not simple. Then there exists a $\mathbb{Q}_p[x, x^{-1}]$ -submodule $0 \neq A \neq \mathbb{L}$ such that $A \cdot \mathbb{L} \subseteq A$. Let $B = A \cap L$. Note that $\bar{B} = (B + \Gamma(L))/\Gamma(L)$ is an ideal of $L/\Gamma(L)$ and also an S -submodule. Hence we can apply the inductive hypothesis. Indeed, from the construction of B we see that the ranks of B and L/B are less than r and the x -ranks of B (by Lemma 2.3) and L/B (as a quotient of L) are at most m . \square

3 p -Groups case

We will begin this section by constructing of certain Lie rings associated with uniform powerful groups. The idea of this construction is taking from [8], but in the present form it is suggested by E. Khukhro. Recall that a finite p -group Q is called **powerful** if $Q/Q^{\mathbf{p}}$ is abelian, where $\mathbf{p} = 4$ if $p = 2$ and $\mathbf{p} = p$ if p is odd, and that a powerful p -group P is called **uniform** if the rank of P^{p^i} does not depend on i , as long as $P^{p^i} \neq 1$. We suggest to the reader the books [1, 4] for detailed properties of these groups.

In this section we will use the following notation. Let G be a uniform powerful p -group and p^n the exponent of G . Define $G_i = G^{p^i}$. We write n in the form $n = 4e + f$, where $f \in [0, 3]$. Put $L = G_e/G_{2e}$ and for each integer $k \in [0, 2]$ let $\pi_k: G_{ke}/G_{(k+1)e} \rightarrow G_{(k+1)e}/G_{(k+2)e}$ be the application defined in the following way: if $t \in G_{ke}$ then $\pi_k(tG_{(k+1)e}) = t^{p^e}G_{(k+2)e}$. It is known that these application is well defined and, moreover, for $k \geq 1$ the application π_k is an isomorphism of groups.

The group L is abelian and we will write the group operation of L in additive form. If $a, b \in G_e$ define

$$[aG_{2e}, bG_{2e}]_L = \pi_1^{-1}([a, b]G_{3e}).$$

Lemma 3.1. *With this definition of brackets, $(L, [,]_L, +)$ is a Lie ring.*

Proof. We will prove that the Jacobi condition holds. The rest of axioms of Lie ring can be proved by the same way. Let $a, b, c \in G_e$. Bearing in mind that $[G_e, G_{3e}] \leq G_{4e}$ and using the Hall-Witt identity, we obtain that

$$[a, b, c][b, c, a][c, a, b] \in G_{4e}.$$

Note that if $d \in G_{2e}$ and $c \in G_e$, then

$$\pi_2([\pi_1^{-1}(dG_{3e}), c]G_{3e}) = [d, c]G_{4e}.$$

Hence

$$[\pi_1^{-1}([a, b]G_{3e}), c][\pi_1^{-1}([b, c]G_{3e}), a][\pi_1^{-1}([c, a]G_{3e}), b] \subseteq G_{3e}.$$

Applying π_1^{-1} to the last equality, we obtain the Jacobi identity. \square

If ϕ is an automorphism of G then ϕ acts in the natural way on L and from the definition of the Lie brackets of L follows that ϕ is a Lie ring automorphism of L .

We will say that a subgroup N of a finite p -group Q is **powerfully embedded in Q** if $N^p \leq [N, Q]$. It is known that if M and N are powerfully embedded in Q , then so are MN , M^p and $[M, N]$. The next lemma which was proved in [9] is very important for future application of our construction.

Lemma 3.2. *If M and N are powerfully embedded subgroups in a finite p -group Q , then $[M^{p^i}, N^{p^j}] = [M, N]^{p^{i+j}}$.*

Lemma 3.3. *If T is powerfully embedded subgroup in G , then $\bar{T} = \pi_1(TG_e)$ is an ideal of L and $[\bar{T}, \bar{T}]_L = \pi_1([T, T]G_e)$.*

Proof. Since TG_e is powerful, $\bar{T} = (TG_e)^{p^e}/G_{2e}$ and so it is a subgroup of L . Let us prove that it is also an ideal of L . Using that $(TG_e)^{p^e}$ is powerfully embedded in G , we obtain that $[(TG_e)^{p^e}, G_e] = [TG_e, G]^{p^{2e}}$. Hence $[\bar{T}, L]_L = [TG_e, G]^{p^e}G_{2e}/G_{2e} \subseteq \bar{T}$.

Now, we will prove the second part of the lemma. We have that

$$[(TG_e)^{p^e}, (TG_e)^{p^e}]G_{3e} = [(TG_e), (TG_e)]^{p^{2e}}G_{3e} = ([T, T]G_e)^{p^{2e}}.$$

Therefore $[\bar{T}, \bar{T}]_L = \pi_1([T, T]G_e)$. \square

Corollary 3.4. *Let T be a powerfully embedded subgroup in G and consider $\bar{T} = \pi_1(TG_e)$ as an ideal of L . Then $\bar{T}^{(d)} = \pi_1(T^{(d)}G_e)$.*

For each integer nonnegative i , we define $L_i = G_{i+e}G_{2e}/G_{2e}$. Then by Lemma 3.3 we know that L_i is an ideal of L .

Lemma 3.5. *Let d be the derived length of L_i (as a Lie ring). Then the derived length of G_i is $\leq d + 2$.*

Proof. By the previous corollary, $G_i^{(d)} \leq G_e$. Now, it is easy to see that $G_i^{(d+2)} = \{1\}$. \square

Proof of Theorem 1.1 for finite p -groups. Let G be a finite p -group of rank r admitting an automorphism ϕ with m fixed points. The case $m = 1$ was considered in [8, ch.4], so we may assume that $m \neq 1$. By [7] there exists a characteristic powerful subgroup Q such that $|G : Q| \leq p^{r \lceil \log_2 r \rceil}$. Hence we can suppose that G is powerful. Since the rank of G is r , there exist $k \leq r$ and characteristic subgroups $\{1\} = G_0 \subset G_1 \subset \dots \subset G_k = G$ such that $H_i = G_i/G_{i-1}, i = 1, \dots, k$, are uniform powerful p -groups. Note that $C_{H_i}(\phi) \leq m$. By the above construction we can associate with each group H_i a Lie ring $L = L(i)$. Moreover, ϕ can be considered as a Lie automorphism of L and we have that $C_L(\phi) \leq m$. By Theorem 2.1 there are functions $t_i = t_i(r, m)$ and $s_i = s_i(r)$ such that $L_f^{(s)} = \{0\}$. Hence by Lemma 3.5, we obtain that $(H_i^{p^{t_i}})^{(s_i+2)} = \{1\}$ and so $(G_i^{p^{t_i}})^{(s_i+2)} \subseteq G_{i-1}$. Hence by Lemma 3.2 we can find $f = f(m, r)$ and $g = g(r)$ such that $(G^{p^f})^{(g)} = \{1\}$. Since G is powerful, the index of G^{p^f} in G is at most p^{fr} . \square

4 Final remarks

Proof of Theorem 1.1. First suppose that G is a nilpotent group. Then G is the direct product of its Sylow p_i -subgroups G_i . Decompose m as $m = \prod p_i^{m_i}$. If $m_i = 0$ then by [8], the derived length of G_i is bounded by a some function which depends only on r . In this case we put $H_i = G_i$. If $m_i \neq 0$, then by the previous section, there exist a characteristic subgroup H_i of G_i of (m_i, r) -bounded index and r -bounded derived length. Then $H = \prod H_i$ is a characteristic subgroup of G of (m, r) -bounded index and r -bounded derived length.

Suppose now that G is a soluble group. We follow the argument from [3]. Let $q \in \pi(G)$ be a prime divisor of $|G|$. Define by $O_{q'}(G)$ the maximal normal q' -subgroup of G and by $O_{qq'}(G)$ the inverse image of the maximal normal q -subgroup Q of the quotient $G/O_{q'}(G)$. It is well-known that the action of $G/O_{qq'}(G)$ by conjugation on $Q/\Phi(Q)$ is faithful. Then the group $G/O_{qq'}(G)$ is isomorphic to a soluble subgroup of $\text{GL}_d(q)$, where d is the number of generators of Q . Since $d \leq r$, using the Kolchin-Malcev theorem [11, Theorem 3.6], we obtain that the derived length of $G/O_{qq'}(G)$ is r -bounded. Hence G/F , where $F = \cap_{q \in \pi(G)} O_{qq'}(G)$ is the Fitting subgroup of G , is soluble of r -bounded derived length. Since F is nilpotent, there exists a characteristic subgroup N of F of (m, r) -bounded index and r -bounded derived length. Put $H = C_G(F/N)$. The index of H is (m, r) -bounded, because G/H acts as a group of automorphisms of F/N and the order of F/N is (r, m) -bounded. Also the derived length of H is r -bounded, because the derived lengths of $H/(H \cap F) \cong HF/F$ and $H \cap N$ are r -bounded and $[H \cap F, H \cap F] \leq (H \cap N)$. Then H satisfies the necessary conditions.

Now, if G is an arbitrary finite group of rank r admitting an automorphism with m fixed points, then using the classification of the finite simple groups, it was proved in [8, Proposition 3.2] that G has a characteristic soluble subgroup of (r, m) -bounded index. Therefore this case follows from the previous paragraph.

□

From [8, Ch.5] we can deduce that if G is a finite p -group of rank r admitting a p' -automorphism ϕ with p^m fixed points then the derived length of G is (m, r) -bounded. This result suggests to us the following conjecture.

Conjecture 1. Let G be a finite p -group of rank r admitting an automorphism ϕ with p^m fixed points. Then the derived length of G is (m, r) -bounded. Moreover there are functions $f = f(m, r)$ and $d = d(r)$ such that G has a subgroup of index at most p^f and derived length at most d .

In [2] it was proved that there are functions $f(p, m, n)$ and $h(m)$ such that any finite p -group G with an automorphism of order p^n , whose centralizer has p^m points, has a subgroup of derived length $\leq h(m)$ and index $\leq f(p, m, n)$. Note that in this situation the rank of G is also (p, m, n) -bounded. Therefore, we pose the following problem.

Conjecture 2. Let G be a finite p -group of rank r admitting a p -automorphism ϕ with p^m fixed points. Then there are $f = f(p, m, r)$ and $d = d(m)$ such that G has a subgroup of index at most f and derived length at most d .

References

- [1] J. Dixon, M. du Sautoy, A. Mann, y D. Segal, *Analytic pro- p groups*, 2nd ed., Cambridge University Press, Cambridge, 1999.
- [2] A. JAIKIN ZAPIRAIN, On the almost regular automorphisms of finite p -groups, *Advances in Mathematics* **153**(2000), 391–402.
- [3] E. I. KHUKHRO, Almost regular automorphisms of finite groups of bounded rank. *Sibirsk. Mat. Zh.*, **37** (1996), no. 6, 1407–1412; English transl. *Siberian Math. J.*, **37** (1996), no. 6, 1237–1241.
- [4] E. I. KHUKHRO, *p -Automorphisms of Finite p -groups*, Cambridge University Press, 1998.
- [5] *The Kourovka notebook: unsolved problems in group theory*, E. I. Khukhro (ed.), V. D. Mazurov (ed.), Novosibirsk, 14th ed., 1999.
- [6] V. A. KREKNIN, The solubility of the Lie algebras with a regular automorphism, *Sibirsk. Math. Zh.* **8** (1967), 715–716; English. transl., *Siberian Math. J.* **8** (1968), 536–537.
- [7] A. LUBOTZKY AND A. MANN, Powerful p -groups. I. Finite groups., *J. Algebra*, **105** (1987), 484-505.
- [8] A. SHALEV, Automorphisms of finite groups of bounded rank, *Israel J. Math.*, **82**(1993), 395-404.

- [9] A. SHALEV, On almost fixed point free automorphisms, *J. Algebra*, **157**(1993), 271–282.
- [10] A. SHALEV, Finite p -groups, Collection: Finite and locally finite groups (Istanbul, 1994), NATO Adv. Sci. Inst. Ser. C Math. Phys. Sci., **471** (1995), 401–450.
- [11] B. A. F. WEHRFRITZ, *Infinite Linear Groups*, Springer-Verlag, 1973.