

# CLOSE LATTICE POINTS ON CIRCLES

JAVIER CILLERUELO AND ANDREW GRANVILLE

ABSTRACT. We classify the sets of four lattice points that all lie on a short arc of a circle which has its center at the origin; specifically on arcs of length  $tR^{1/3}$  on a circle of radius  $R$ , for any given  $t > 0$ . In particular we prove that any arc of length  $(40 + \frac{40}{3}\sqrt{10})^{1/3} R^{1/3}$  on a circle of radius  $R$ , with  $R > \sqrt{65}$ , contains at most three lattice points, whereas we give an explicit infinite family of 4-tuples of lattice points,  $(\nu_{1,n}, \nu_{2,n}, \nu_{3,n}, \nu_{4,n})$ , each of which lies on an arc of length  $(40 + \frac{40}{3}\sqrt{10})^{1/3} R_n^{1/3} + o(1)$  on a circle of radius  $R_n$ .

## 1. INTRODUCTION

How many lattice points  $(x, y) \in \mathbb{Z}^2$  can be on a “small” arc of the circle  $x^2 + y^2 = R^2$ ?<sup>1</sup> A. Córdoba and the first author [3] proved that for every  $\epsilon > 0$  the number of lattice points on an arc of length  $R^{\frac{1}{2}-\epsilon}$  is bounded uniformly in  $R$ . More precisely they proved (see also [4] and [6]):

**Theorem 1.1.** *For any integer  $k \geq 1$ , an arc of length  $\sqrt{2}R^{\frac{1}{2}-\frac{1}{4\lfloor k/2 \rfloor + 2}}$  on a circle of radius  $R$  centered at the origin contains no more than  $k$  lattice points.*

This result cannot be improved for  $k = 1$  since the circles  $x^2 + y^2 = 2n^2 + 2n + 1$  contain two lattice points,  $(n, n + 1)$  and  $(n + 1, n)$ , on an arc of length  $\sqrt{2} + o(1)$ .

Theorem 1.1 for  $k = 2$  was first proved by Schinzel, and then used by Zygmund [10] to prove a result about spherical summability of Fourier series in two dimensions. In [2] the first author gave a best possible version of Schinzel’s result (which we will prove more easily in section 2).

**Theorem 1.2.** *An arc of length  $(16R)^{\frac{1}{3}}$  on a circle of radius  $R$  centered at the origin contains no more than two lattice points.*

This result cannot be improved since the circles  $x^2 + y^2 = R_n^2 := 16n^6 + 4n^4 + 4n^2 + 1$  contain three lattice points,  $(4n^3 - 1, 2n^2 + 2n)$ ,  $(4n^3, 2n^2 + 1)$  and  $(4n^3 + 1, 2n^2 - 2n)$ , on an arc of length  $(16R_n)^{\frac{1}{3}} + o_n(1)$ .

Let  $[\nu] = (\nu_1, \dots, \nu_k)$  denote a  $k$ -tuple of lattice points lying on the same circle of radius  $R = R_{[\nu]}$  centered at the origin, and  $\text{Arc}[\nu] = \text{Arc}(\nu_1, \dots, \nu_k)$  the length of the shortest arc containing  $\nu_1, \dots, \nu_k$ .

---

1991 *Mathematics Subject Classification.* 2000 *Mathematics Subject Classification:*11N36.

Le deuxième auteur est partiellement soutenu par une bourse du Conseil de recherches en sciences naturelles et en génie du Canada. He would like to thank the first author for his frequent hospitality, as well that of the Universidad Autónoma de Madrid, over the years in which we worked together on this stubborn problem.

<sup>1</sup>If there are points with integer coordinates on the circle  $x^2 + y^2 = R^2$  then  $R^2$  must be an integer. Henceforth we shall assume this, whether we state it or not.

The next result shows that we cannot improve the constant  $(16)^{1/3}$  if we omit the examples above.

**Theorem 1.3.** *The set  $\{\text{Arc}[\nu]R_{[\nu]}^{-1/3}, [\nu] = (\nu_1, \nu_2, \nu_3)\}$  is dense in  $[(16)^{1/3}, +\infty)$ .*

Since we have sharp versions of Theorem 1.1 for  $k = 1$  and  $2$ , we focus in this paper on giving a sharp version of Theorem 1.1 for  $k = 3$ . We begin by showing that the exponent given in Theorem 1.1 is best possible for  $k = 3$ , by exhibiting infinitely many circles  $x^2 + y^2 = R^2$  with four lattice points in an arc of length  $\ll R^{1/3}$ : The Fibonacci numbers are defined by  $F_0 = 0$ ,  $F_1 = 1$  and  $F_{n+2} = F_{n+1} + F_n$  for all  $n \geq 0$ . The circles  $x^2 + y^2 = R_n^2 := \frac{5}{2}F_{2n-1}F_{2n+1}F_{2n+3}$  contain the four lattice points  $\frac{1}{2}(F_{3n+3}, F_{3n}) + (-1)^n z_j$  for  $j = 1, 2, 3, 4$ , where

$$z_1 = 2(-F_{n-1}, F_{n+2}), \quad z_2 = (-F_{n-2}, F_{n+1}), \quad z_3 = (F_{n-1}, -F_{n+2}), \quad z_4 = (F_n, -F_{n+3}).$$

The chord length between  $z_1$  and  $z_4$  is  $\sqrt{10F_{2n+3}}$ , implying that the arc containing all four lattice points has length

$$2R_n \arcsin\left(\frac{\sqrt{10F_{2n+3}}}{2R_n}\right) = 20^{\frac{1}{3}} \left(\frac{1+\sqrt{5}}{2}\right) R_n^{1/3} + \frac{2\sqrt{5}}{3R_n} + O\left(\frac{1}{R_n^{7/3}}\right).$$

In fact the arc length can be shown to be  $> 20^{\frac{1}{3}} \left(\frac{1+\sqrt{5}}{2}\right) R_n^{1/3}$ .

We see here a family  $\mathcal{F} = \{[\nu]_n = (\nu_{1,n}, \nu_{2,n}, \nu_{3,n}, \nu_{4,n}), n \in \mathbb{N}\}$  of 4-tuples of lattice points, lying on circles centered at the origin, with

$$\text{Arc}([\nu]_n) \sim C_{\mathcal{F}} R_n^{1/3} \quad \text{as } n \rightarrow \infty.$$

The main result of this paper is that 4-tuples of lattice points which lie on a short arc of a circle centered at the origin, belong to such a family, and that  $\mathcal{F}(t) = \{\mathcal{F}, C_{\mathcal{F}} \leq t\}$  is a finite set for any given  $t$  (which is rather different from the  $k = 2$  case, as given in Theorem 1.3).

**Theorem 1.4.** *For any  $t > 0$  any arc on the circle  $x^2 + y^2 = R^2$ , of length less than  $tR^{1/3}$  with  $R > 2^{-18}t^{15}$ , contains at most three lattice points except for the families  $\mathcal{F} \in \mathcal{F}(t)$ .*

Therefore, in contrast to Theorem 1.3, we deduce the following.

**Corollary 1.** *The set  $\{\text{Arc}[\nu]R_{[\nu]}^{-1/3}, [\nu] = (\nu_1, \nu_2, \nu_3, \nu_4)\}$  has only finitely many accumulation points in any interval  $[0, t)$ , where  $t \in \mathbb{R}^+$ .*

We order the families  $\mathcal{F}_1, \mathcal{F}_2, \dots$  so that  $C_{\mathcal{F}_1} \leq C_{\mathcal{F}_2} \leq \dots$ . For fixed  $t$  we can explicitly determine  $\mathcal{F}(t)$  (as described in section 8); indeed, in the table there we describe all seven families belonging to  $\mathcal{F}(5)$ . We found that  $C_{\mathcal{F}_1} = (40 + \frac{40}{3}\sqrt{10})^{1/3}$ , and then  $C_{\mathcal{F}_2} = 20^{\frac{1}{3}} \left(\frac{1+\sqrt{5}}{2}\right)$  where  $\mathcal{F}_2$  is the family given above. We deduce the following

**Corollary 2.** *An arc of the circle  $x^2 + y^2 = R^2$ , with  $R > \sqrt{65}$ , of length  $< (40 + \frac{40}{3}\sqrt{10})^{1/3} R^{1/3}$  contains at most three lattice points. On the contrary, there are infinitely many circles  $x^2 + y^2 = R_n^2$  containing four lattice points in arcs on length  $(40 + \frac{40}{3}\sqrt{10})^{1/3} R_n^{1/3} + o(1)$ .*

Note that  $16^{1/3} = 2.5198\dots < (40 + \frac{40}{3}\sqrt{10})^{1/3} = 4.347\dots < 20^{\frac{1}{3}} \left(\frac{1+\sqrt{5}}{2}\right) = 4.3920\dots$

The coordinates of the 4-tuples in each family  $\mathcal{F}$  grow exponentially. This implies that for each fixed  $t > 0$  there exists a constant  $B_t$  (which will be described in section 9) such that there are  $\sim B_t \log x$  4-tuples of lattice points which lie on an arc of length  $tR^{1/3}$  of a circle of radius  $R$  centered at the origin, where  $R \leq x$ .

## 2. THREE LATTICE POINTS

We give here the proof of several results that were discussed in the introduction. Our new proof of Theorem 1.2 is somewhat simpler than that in [2].

*Proof of Theorem 1.2.* Suppose that  $\nu_1, \nu_2, \nu_3$  are three lattice points, in order, on a circle of radius  $R$  so that

$$|\nu_1 - \nu_2||\nu_2 - \nu_3||\nu_1 - \nu_3| < \text{Arc}(\nu_1, \nu_2)\text{Arc}(\nu_2, \nu_3)\text{Arc}(\nu_1, \nu_3) \leq \frac{1}{4}\text{Arc}(\nu_1, \nu_3)^3.$$

A theorem attributed to Heron of Alexandria states that if  $\Delta$  is the area of the triangle with sides  $a, b, c$ , and  $R$  is the radius of the circle going through the vertices of the triangle, then  $abc = 4\Delta R$ . Applying this to the triangle with vertices  $\nu_1, \nu_2, \nu_3$  we have that  $|\nu_1 - \nu_2||\nu_2 - \nu_3||\nu_1 - \nu_3| = 4\Delta R$ .

It should be noted that any triangle with integer vertices has area  $\geq 1/2$  so, a priori,  $\Delta \geq 1/2$ . However, we can do better than this: Since  $\nu_1, \nu_2, \nu_3$  lie on the same circle, an easy parity argument implies that the co-ordinates of two of these lattice points, say  $\nu_i \neq \nu_j$ , have the same parity, and so  $\frac{1}{2}(\nu_i + \nu_j)$  is also an integer lattice point. Therefore the triangle  $\nu_1, \nu_2, \nu_3$  is the disjoint union of two triangles with integer coordinates, which implies that  $\Delta \geq 1$ . The result follows.  $\square$

The second author posed a weak version of Theorem 1.2 as problem A5 on the 2000 Putnam examination; about 45 contestants had the wherewithal to provide a solution somewhat like that above.

Henceforth we identify the lattice point  $(x, y) \in \mathbb{Z}^2$  with the Gaussian integer  $x + iy$ .

*Proof of Theorem 1.3.* Let  $C \geq (16)^{1/3}$  and  $\alpha$  satisfying  $(1 + \alpha) \left(\frac{4}{\alpha + \alpha^2}\right)^{1/3} = C$ . Take  $p$  and  $q$  to be distinct large primes for which  $n_2 \sim \alpha n_1$  where  $n_1 = 2p$  and  $n_2 = q$ . Now, take  $m_1$  to be an odd integer and  $m_2$  to be an even integer, much larger than  $n_1$  and  $n_2$ , such that  $m_1 n_2 - m_2 n_1 = \pm 1$ . Finally take  $n_3 = \frac{1}{2}(n_1 + n_2 + m_1 + m_2)$  and  $m_3 = \frac{1}{2}(n_1 + n_2 - m_1 - m_2)$ . We write  $\mu_j + im_j$ ,  $j = 1, 2, 3$  and consider

$$\nu_1 = \mu_1 \bar{\mu}_2 \mu_3 \quad \nu_2 = i \bar{\mu}_1 \bar{\mu}_2 \bar{\mu}_3, \quad \nu_3 = \bar{\mu}_1 \mu_2 \mu_3.$$

Notice that  $|\mu_1| \sim m_1$ ,  $|\mu_2| \sim m_1 \alpha$  and  $|\mu_3| \sim m_1(1 + \alpha)/\sqrt{2}$ , so that

$$R^{1/3} = |\nu_j|^{1/3} \sim ((\alpha + \alpha^2)/\sqrt{2})^{1/3} m_1.$$

Now  $|\nu_3 - \nu_1|R^{-1/3} = |\mu_3||\mu_1 \bar{\mu}_2 - \bar{\mu}_1 \mu_2|R^{-1/3} = 2|\mu_3|R^{-1/3} \sim (1 + \alpha) \left(\frac{4}{\alpha + \alpha^2}\right)^{1/3} = C$ , and similarly both  $|\nu_3 - \nu_2|R^{-1/3} \sim \left(\frac{4}{\alpha + \alpha^2}\right)^{1/3}$  and  $|\nu_2 - \nu_1|R^{-1/3} \sim \alpha \left(\frac{4}{\alpha + \alpha^2}\right)^{1/3}$ .  $\square$

## 3. THE CONSTRUCTION OF THE FAMILIES OF 4-TUPLES OF LATTICE POINTS

If  $\mathbf{g} = \gcd(\nu_1, \dots, \nu_k)$  then  $\frac{\text{Arc}[\nu]}{R} = \frac{\text{Arc}[\nu/\mathbf{g}]}{R/|\mathbf{g}|}$  so that  $\text{Arc}[\nu] = |\mathbf{g}|\text{Arc}[\nu/\mathbf{g}]$ , so we can reduce our study to primitive 4-tuples of lattice points where  $[\nu]$  is *primitive* if  $\gcd(\nu_1, \nu_2, \nu_3, \nu_4) = 1$ .

We therefore consider primitive 4-tuples of lattice points  $[\nu] = (\nu_1, \nu_2, \nu_3, \nu_4)$  which all lie on the same circle centered at the origin, say  $x^2 + y^2 = R^2$  and we assume that

$$(3.1) \quad \sigma_n := \nu_1 + \nu_2 + \nu_3 + \nu_4 \neq 0.$$

(Note that if  $\sigma = 0$  then the  $\nu_i$  cannot all lie on the same half circle, and hence  $\text{Arc}[\nu] \geq \pi R$ ; we shall have more to say about this case at the start of section 5.) Next define

$$\omega_{[\nu]} = \left( \frac{\nu_1 \nu_2 \nu_3 \nu_4}{|\nu_1 \nu_2 \nu_3 \nu_4|} \right)^{\frac{1}{4}} = \frac{(\nu_1 \nu_2 \nu_3 \nu_4)^{1/4}}{R} \quad \text{so that} \quad -\pi/4 < \text{Arg}(\sigma_{[\nu]} \overline{\omega_{[\nu]}}) \leq \pi/4.$$

Let  $\Psi_{[\nu]} := \text{Arg}(\sigma_{[\nu]} \overline{\omega_{[\nu]}})$ , so that  $-1 < \tan(\Psi_{[\nu]}) \leq 1$  and  $\cos(\Psi_{[\nu]}) > 0$ .

Let  $Q = Q_{[\nu]}$  be the smallest positive integer for which  $\sqrt{Q}\omega^2 \in \mathbb{Z}[i]$  (we will prove that  $Q$  exists in section 4). If  $Q_{[\nu]}$  is a square then  $[\nu]$  is *degenerate*, a simple case that we will examine in section 5. Typically  $Q_{[\nu]}$  is not a square, that is  $[\nu]$  is non-degenerate, in which case we select the smallest possible positive integers  $p$  and  $q$  for which<sup>2</sup>

$$p^2 - q^2 Q = \epsilon = \pm 1,$$

and we write  $\alpha := p + q\sqrt{Q}$  and  $\beta := p - q\sqrt{Q}$ .

For a given  $[\nu]$ , we define the complex numbers<sup>3</sup>

$$\omega_1 = \frac{(\nu_1 \overline{\nu}_2 \overline{\nu}_3 \nu_4)^{1/4}}{R}, \quad \omega_2 = \frac{(\overline{\nu}_1 \nu_2 \overline{\nu}_3 \nu_4)^{1/4}}{R} \quad \text{and} \quad \omega_3 = \frac{(\overline{\nu}_1 \overline{\nu}_2 \nu_3 \nu_4)^{1/4}}{R}.$$

For each integer  $n$  we define

$$(3.2) \quad \omega_{i,n} = \alpha^n \frac{\omega_i + \overline{\omega}_i}{2} + \beta^n \frac{\omega_i - \overline{\omega}_i}{2}, \quad i = 1, 2, 3.$$

and then a sequence of 4-tuples of lattice points  $\{[\nu]_n = (\nu_{1,n}, \nu_{2,n}, \nu_{3,n}, \nu_{4,n}), n \in \mathbb{Z}\}$  by

$$(3.3) \quad \begin{aligned} \nu_{1,n} &= R\omega \omega_{1,n} \overline{\omega}_{2,n} \overline{\omega}_{3,n} \\ \nu_{2,n} &= R\omega \overline{\omega}_{1,n} \omega_{2,n} \overline{\omega}_{3,n} \\ \nu_{3,n} &= R\omega \overline{\omega}_{1,n} \overline{\omega}_{2,n} \omega_{3,n} \\ \nu_{4,n} &= R\omega \omega_{1,n} \omega_{2,n} \omega_{3,n} \end{aligned}$$

We immediately deduce that the lattice points  $\nu_{j,n}$ ,  $j = 1, 2, 3, 4$ , all lie on the same circle, and that  $\omega_{[\nu]_n}^4 = \omega^4$ . Multiplying out the terms in this definition we obtain

$$(3.4) \quad \begin{aligned} \nu_{j,n} &= \alpha^{3n} \frac{\sigma + \omega^2 \overline{\sigma}}{8} + (\epsilon\alpha)^n \left( \frac{\nu_j - \omega^2 \overline{\nu}_j}{2} - \frac{\sigma - \omega^2 \overline{\sigma}}{8} \right) \\ &\quad + \beta^{3n} \frac{\sigma - \omega^2 \overline{\sigma}}{8} + (\epsilon\beta)^n \left( \frac{\nu_j + \omega^2 \overline{\nu}_j}{2} - \frac{\sigma + \omega^2 \overline{\sigma}}{8} \right) \end{aligned}$$

so that

$$(3.5) \quad \begin{aligned} \nu_{j,n} \overline{\omega} &= \frac{\alpha^{3n}}{4} \text{Re}(\sigma \overline{\omega}) + i(\epsilon\alpha)^n \left( \text{Im}(\nu_j \overline{\omega}) - \frac{\text{Im}(\sigma \overline{\omega})}{4} \right) \\ &\quad + i \frac{\beta^{3n}}{4} \text{Im}(\sigma \overline{\omega}) + (\epsilon\beta)^n \left( \text{Re}(\nu_j \overline{\omega}) - \frac{\text{Re}(\sigma \overline{\omega})}{4} \right) \end{aligned}$$

for  $j = 1, 2, 3, 4$ . We deduce that  $\nu_{j,n} = \omega R_n + O(R_n^{1/3})$  as  $\alpha > 1 > |\beta|$ , and that

$$\sigma_n := \sum_{j=1}^4 \nu_{j,n} = \alpha^{3n} \frac{\sigma + \omega^2 \overline{\sigma}}{2} + \beta^{3n} \frac{\sigma - \omega^2 \overline{\sigma}}{2} = \omega \left( \alpha^{3n} \text{Re}(\sigma \overline{\omega}) + i\beta^{3n} \text{Im}(\sigma \overline{\omega}) \right).$$

<sup>2</sup>But see remark 1

<sup>3</sup>There is some ambiguity here, in that these quantities are well-defined only up to a fourth root of unity. Our protocol is to make a choice for the value of each  $\nu_j^{1/4}/R$  (out of the four possibilities) so as to validate the choice of fourth root of unity in the definition of  $\omega = \omega_{[\nu]}$ , and then to use this same value for  $\nu_j^{1/4}/R$  consistently throughout these definitions.

Now  $|\tan(\text{Arg}(\sigma_n \bar{\omega}))| = \left| \frac{\beta^{3n} \text{Im}(\sigma \bar{\omega})}{\alpha^{3n} \text{Re}(\sigma \bar{\omega})} \right| = |\beta^{6n} \tan \Psi_{[\nu]}| < 1$  as  $|\beta| < 1$  if  $n \geq 0$ , and also  $\cos(\text{Arg}(\sigma_n \bar{\omega})) > 0$  as  $\alpha > 0$  and  $\text{Re}(\sigma \bar{\omega}) > 0$ , which implies that  $\omega_{[\nu]_n} = \omega$  (since we already know that  $\omega_{[\nu]_n}^4 = \omega^4$ ). In other words,  $\omega$  is an invariant of the family, and so  $Q = Q_{\mathcal{F}}$  is also.

With a formula like (3.4) it is evident that one can express the  $\nu_{j,n}$  in terms of a recurrence: For  $n \geq 0$  we have

$$(3.6) \quad \nu_{j,n+1} = pq \left( qQ\sigma_n + p\sqrt{Q}\omega^2\bar{\sigma}_n \right) + \epsilon \left( p\nu_{j,n} - q\sqrt{Q}\omega^2\bar{\nu}_{j,n} \right).$$

We deduce that each  $\nu_{j,n} \in \mathbb{Z}[i]$  by induction on  $n \geq 0$ , since  $\sqrt{Q}\omega^2 \in \mathbb{Z}[i]$ . This completes the proof that each  $[\nu]_n$  with  $n \geq 0$  gives rise to a 4-tuple of lattice points on a circle centered at the origin.

We can re-express (3.6) in the more friendly looking form

$$(3.7) \quad \nu_{j,n} = aG_{3n} + bG_{3n+1} + \epsilon^n (a_j G_n + b_j G_{n+1}) \quad \text{for all } n \geq 0$$

with  $b = \sigma/4$ ,  $a = q\sqrt{Q}\omega^2\bar{b} - pb$ , and  $b_j = \nu_j - \sigma/4$ ,  $a_j = q\sqrt{Q}\omega^2\bar{b}_j - pb_j$  for each  $j$ , where the recurrence sequence  $\{G_n : n \geq 0\}$  is defined by

$$(3.8) \quad G_0 = 0, \quad G_1 = 1, \quad \text{and} \quad G_n = 2pG_{n-1} - \epsilon G_{n-2} \quad \text{for all } n \geq 2.$$

**Remark 1.** *This formula can be used to show that one can obtain Gaussian integers  $\nu_{j,n}$  even when  $p, q \in \mathbb{Z} + \frac{1}{2}$  (instead of in  $\mathbb{Z}$ ); we take these semi-integer values for  $p = p_{[\nu]}$ ,  $q = q_{[\nu]}$  whenever possible. For example if  $[\nu] = (1-2i, 2-i, 2+i, -1+2i)$  then  $Q = 5$  and we can take  $p = q = 1/2$  (as we now verify): The sequence  $G_n$  in (3.8) is the Fibonacci sequence  $F_n$ , and we have  $a_1 = -1+3i$ ,  $b_1 = -2i$ ,  $a_2 = 2i$ ,  $b_2 = 1-i$ ,  $a_3 = 1-i$ ,  $b_3 = 1+i$ ,  $a_4 = -4i$ ,  $b_4 = -2+2i$ , so that (3.7) gives*

$$\nu_{j,n} = \frac{1+i}{2} F_{3n} + F_{3n+1} + a_j F_n + b_j F_{n+1}.$$

Hence  $\nu_{j,n}$  is always a Gaussian integer since  $F_{3n}$  is always an even rational integer (and we obtain the example given in the introduction).

What about  $n < 0$ ? The above proof is easily modified to work for all negative  $n$  except the requirement that  $|\beta^{6n} \tan \Psi_{[\nu]}| < 1$ . Thus we select  $n_0$  to be the smallest integer for which  $|\beta^{6n} \tan \Psi_{[\nu]}| < 1$  and replace  $[\nu]$  by  $[\nu]_{n_0}$ . We call this an *initial 4-tuple*, and then define the family

$$\mathcal{F}([\nu]) = \{(\nu_{1,n}, \nu_{2,n}, \nu_{3,n}, \nu_{4,n}), \quad n \geq 0\}.$$

We extend this definition to any  $[\nu]$  by defining  $\mathcal{F}([\nu]) = \mathcal{F}([\nu]_{n_0})$  for  $[\nu] \in \mathcal{F}([\nu]_{n_0})$ . If  $[\nu]$  is an initial 4-tuple then  $[\nu]_{-1}$  is an initial 4-tuple of a different family, the *dual family*, which we denote by  $\hat{\mathcal{F}} := \mathcal{F}([\nu]_{-1})$ .

By (3.5) we see that  $R_n \sim \frac{\alpha^{3n}}{4} \text{Re}(\sigma \bar{\omega})$  and  $|\nu_{j,n} - \nu_{k,n}| \sim \alpha^n |\text{Im}((\nu_j - \nu_k) \bar{\omega})|$ , as  $\alpha > 1 > |\beta|$ , from which we deduce that

$$C_{\mathcal{F}([\nu])} = \max_{1 \leq j < k \leq 4} \frac{|2\text{Im}((\nu_j - \nu_k) \bar{\omega})|}{|2\text{Re}(\sigma \bar{\omega})|^{1/3}},$$

so that

$$(3.9) \quad \text{Arc}([\nu]_n) \sim C_{\mathcal{F}} R_n^{1/3} \quad \text{as } n \rightarrow \infty.$$

Similarly

$$C_{\hat{\mathcal{F}}} = \max_{1 \leq j < k \leq 4} \frac{|2\text{Re}((\nu_j - \nu_k) \bar{\omega})|}{|2\text{Im}(\sigma \bar{\omega})|^{1/3}},$$

and note that  $C_{\hat{\mathcal{F}}}([\nu]) = C_{\mathcal{F}}(i[\nu])$ .

#### 4. PROPERTIES OF $Q$

**Lemma 4.1.** *There exists a positive integer  $Q$ , not divisible by 4, for which  $\sqrt{Q}\omega^2 \in \mathbb{Z}[i]$ . In fact if odd prime  $p$  divides  $Q$  then  $p \equiv 1 \pmod{4}$ . Moreover  $Q/(2, Q)$  divides  $R^2$ .*

*Proof.* Let  $\gamma_i$  be the exact power of prime ideal  $\mathfrak{p}$  of norm  $p$  which divides  $\nu_i$ ,  $i = 1, 2, 3, 4$ , say with  $\gamma_1 \geq \gamma_2 \geq \gamma_3 \geq \gamma_4$ . Since  $[\nu]$  is primitive we know that  $p \neq 2$ ,  $\gamma_4 = 0$  and  $p^{\gamma_1}$  is the exact power of  $p$  dividing  $R^2$ , so that  $\gamma_1 - \gamma_i$  is the exact power of prime ideal  $\bar{\mathfrak{p}}$  which divides  $\nu_i$ . Therefore if  $\gamma = \gamma_1 - \gamma_2 - \gamma_3 - \gamma_4$  then the exact powers of  $\mathfrak{p}$  and  $\bar{\mathfrak{p}}$  dividing  $\omega^4$  are given by  $(\bar{\mathfrak{p}}/\mathfrak{p})^\gamma$  which equals  $(\bar{\mathfrak{p}}^2/p)^\gamma$  if  $\gamma > 0$ , and equals  $(\mathfrak{p}^2/p)^{-\gamma}$  if  $\gamma < 0$ . We see that if  $Q_1$  is the product of these  $p^{|\gamma|}$  then  $Q_1\omega^4 \in u\mathbb{Z}[i]^2$  for some unit  $u$ , since all ideals of  $\mathbb{Z}[i]$  are principal. Taking square roots we see that we can take  $Q = Q_1$  if  $u = \pm 1$ , and  $Q = 2Q_1$  if  $u = \pm i$ , so that  $Q$  is not divisible by 4, and all of its prime factors are norms of elements of  $\mathbb{Z}[i]$  and are thus not  $\equiv 3 \pmod{4}$ .

Finally note that  $|\gamma| = |\gamma_1 - \gamma_2 - \gamma_3| \leq \gamma_1$  so that  $Q_1$  divides  $R^2$ .  $\square$

**Lemma 4.2.** *Let  $\mathfrak{p}$  be a prime ideal in  $\mathbb{Z}[i]$ . If  $[\nu] = (\nu_1, \nu_2, \nu_3, \nu_4)$  is primitive and  $\mathfrak{p}^\alpha$  divides  $\sqrt{Q}\omega^2$ , then  $\mathfrak{p}^\alpha$  divides exactly three of  $\{\nu_1, \nu_2, \nu_3, \nu_4\}$ .*

*Proof.* In the notation of the proof of the previous lemma one finds that the exact power of  $\mathfrak{p}$  which divides  $\sqrt{Q}\omega^2$  is  $\mathfrak{p}^{\max\{0, -\gamma\}}$ , and  $\max\{0, -\gamma\} = \max\{0, \gamma_3 - (\gamma_1 - \gamma_2)\} \leq \gamma_3$ , so the result follows.  $\square$

For given odd integer  $n$  we define

$$r(n) := \min_{\substack{r_1 r_2 r_3 r_4 = n \\ (r_i, r_j) = 1, i \neq j}} \max_{1 \leq i \leq 4} r_i$$

and then let  $r(2^k n) = r(n)$ .

**Lemma 4.3.** *If  $[\nu]$  is a primitive 4-tuple we have  $\text{Arc}[\nu] > (16r(Q_{[\nu]}))^{1/3} R^{1/3}$ .*

*Proof.* Let  $\mathbf{g}_i = \gcd(\nu_j : j \neq i)$  for  $i = 1, 2, 3, 4$ . By the previous lemma we know that  $\sqrt{Q}\omega^2$  divides  $\mathbf{g}_1 \mathbf{g}_2 \mathbf{g}_3 \mathbf{g}_4$ , so that  $Q$  divides  $|\mathbf{g}_1|^2 |\mathbf{g}_2|^2 |\mathbf{g}_3|^2 |\mathbf{g}_4|^2$ . Therefore there exists some  $j$  for which  $|\mathbf{g}_j|^2 \geq r(Q)$ . Suppose that  $j = 4$  here and let  $\mathbf{g} = \mathbf{g}_4$ . Let  $\nu_i = \mathbf{g}\tau_i$  for  $i = 1, 2, 3$  then  $\text{Arc}[\nu] \geq \text{Arc}[\nu_1, \nu_2, \nu_3] = \text{Arc}(\mathbf{g}[\tau]) = |\mathbf{g}| \text{Arc}([\tau]) \geq |\mathbf{g}| (16R_{[\tau]})^{1/3} = |\mathbf{g}|^{2/3} (16R)^{1/3}$  by Theorem 1.2, as  $R_{[\tau]} = R/|\mathbf{g}|$ . The result follows.  $\square$

We deduce the following result from Lemma 4.3 and (3.9).

**Corollary 3.** *If  $\mathcal{F}$  is a non-degenerate family then  $C_{\mathcal{F}} \geq (16r(Q_{\mathcal{F}}))^{1/3}$ .*

#### 5. DEGENERATE 4-TUPLES

If  $Q$  is a square then  $\alpha = \beta = 1$  so that  $\nu_{j,n} = \nu_j$  for all  $n$  and  $j$ , which explains why we call this the “degenerate” case.

In section 2 we noted that if  $\sigma = 0$  then the  $\nu_i$  cannot all lie on the same half circle, implying that  $\text{Arc}[\nu] \geq \pi R$ ; now we will show that if  $[\nu]$  is also primitive then it is degenerate: Since  $\nu_1 + \nu_2 = (-\nu_3) + (-\nu_4)$  where  $|\nu_1| = |\nu_2| = |-\nu_3| = |-\nu_4|$ , we either have  $\nu_1 + \nu_2 = 0$  (in which case  $\nu_3 + \nu_4 = 0$ ), or that the non-zero sum of two vectors,  $\nu_1$  and  $\nu_2$ , of the same length

equals the sum of two other vectors,  $-\nu_3$  and  $-\nu_4$ , of the same length, and it is then easy to show that those two sets of two vectors must be identical. Thus, by re-ordering the indices if necessary, we have  $\nu_1 + \nu_2 = \nu_3 + \nu_4 = 0$ . But  $(\nu_1, \nu_3) = 1$  since  $[\nu]$  is primitive and hence  $\nu_3 = u\bar{\nu}_1$  where  $u = 1, -1, i$  or  $-i$ . Therefore  $\nu_1\nu_2\nu_3\nu_4 = (\nu_1\nu_3)^2 = u^2R^4$  so that  $\omega_{[\nu]}^2 = \pm u$  and therefore  $Q_{[\nu]} = 1$ .

**Lemma 5.1.** *If  $Q_{[\nu]}$  is a square then  $\text{Arc}[\nu] > 2R^{1/2}/Q^{1/8}$ .*

*Proof.* In the argument of Lemma 4.1 we see that if  $Q$  is a square then each  $\gamma$  must be even and  $u = \pm 1$ . Therefore there exists  $\ell \in \mathbb{Z}[i]$  for which  $Q\omega^4 = \pm\ell^4$ , so that  $|\ell| = Q^{1/4}$ . We deduce that  $(\bar{\ell}\nu_1)(\bar{\ell}\nu_2)(\bar{\ell}\nu_3)(\bar{\ell}\nu_4) = \bar{\ell}^4R^4\omega^4 = \pm|\ell|^8R^4/Q = \pm QR^4$ . If  $\pm = +$  then let  $[\nu'] = \bar{\ell}[\nu]$ ; if  $\pm = -$  let  $[\nu'] = (1+i)\bar{\ell}[\nu]$ . Either way we have  $(\omega')^4 = 1$ , so that  $Q' = 1$ .

Now  $R'^2 \geq 5$  else there are exactly four lattice points on our circle so that  $\sigma_{[\nu]} = 0$ . We may assume that  $\text{Arc}[\nu'] < \frac{\pi}{2}R'$ , else  $\text{Arc}[\nu] \geq 2(R')^{1/2}$  as  $R' \geq \sqrt{5}$ . Therefore, using the obvious symmetries (that is, multiplying  $[\nu']$  through by a unit or replacing it with  $[\bar{\nu}']$ ), we can assume that  $-\pi/2 < \varphi_1 < \varphi_2 < \varphi_3 < \varphi_4 < \pi/2$  where  $\nu'_j = R'e^{i\varphi_j} = x_j + iy_j$ ,  $j = 1, 2, 3, 4$ , and we already know that  $\varphi_1 + \varphi_2 + \varphi_3 + \varphi_4 = 0$ . Thus  $\varphi_1 < 0 < \varphi_4$  and suppose that  $|\varphi_1| \geq \varphi_4$ . This implies that  $\varphi_3 > 0$  so that  $y_3 > 0$  and  $x_3 > x_4$ , and thus  $\text{Arc}[\nu'] > 2y_4 = 2\sqrt{R'^2 - x_4^2} \geq 2\sqrt{R'^2 - (x_3 - 1)^2} \geq 2\sqrt{R'^2 - (R' - 1)^2} = 2\sqrt{2R' - 1} > 2^{5/4}(R')^{1/2}$  as  $R' \geq \sqrt{5}$ . Therefore from the remarks at the beginning of section 3 we have

$$\text{Arc}[\nu] = \frac{\text{Arc}[\nu']}{|\ell'|} > \frac{2^{5/4}(R')^{1/2}}{|\ell'|} = \frac{2^{5/4}R^{1/2}}{|\ell'|^{1/2}} \geq \frac{2R^{1/2}}{Q^{1/8}}.$$

□

**Corollary 4.** *If  $\text{Arc}[\nu] < tR^{1/3}$  and  $[\nu]$  is primitive and degenerate then  $R \leq t^{15}/2^{18}$ .*

*Proof.* If  $Q_{[\nu]}$  is an square then  $\text{Arc}[\nu] > \max\{2R^{1/2}/Q^{1/8}, (16r(Q)R)^{1/3}\}$  by Lemmas 4.3 and 5.1. Therefore  $\text{Arc}[\nu] > 2^{6/5}R^{2/5}$  as  $r(Q) \geq Q^{1/4}$ , so  $R < t^{15}/2^{18}$  and the result follows. □

## 6. THE CONSTANT $C_{\mathcal{F}}$ ASSOCIATED TO A FAMILY $\mathcal{F}$

We begin this section by noting, without proof, two technical trigonometric lemmas that will be useful below.

**Lemma 6.1.** *If  $-\pi/2 \leq x_1 < \dots < x_n \leq \pi/2$  then*

$$(6.1) \quad \max_{1 \leq i < j \leq n} |\sin(x_i) - \sin(x_j)| = |\sin(x_1) - \sin(x_n)|.$$

**Lemma 6.2.** *If  $|x| \leq \pi/4$  then*

- i)  $|\sin x| < 1.0106|x| \cdot |\cos x|^{1/3}$ .
- ii)  $|\sin x| < |x| \cdot \left|\frac{\cos x + \cos y}{2}\right|^{1/3}$  whenever  $|y| \leq |x| - 0.137|x|^3$
- iii)  $|\sin x| |\cos x|^{1/3} \geq \max(2|x|/\pi, |x| - |x|^3/3)$ .

The following is the main result in the section.

**Theorem 6.1.** *If  $\text{Arc}[\nu] < \frac{\pi}{2}R_{[\nu]}$  where  $[\nu]$  is primitive and non-degenerate then*

- i)  $\text{Arc}[\nu] > 0.9895C_{\mathcal{F}}R_{[\nu]}^{1/3}$ .
- ii)  $\text{Arc}[\nu] > C_{\mathcal{F}}R_{[\nu]}^{1/3}$  for  $R_{[\nu]} > 0.08C_{\mathcal{F}}^{15/4}$ .
- iii)  $\text{Arc}[\nu] \leq C_{\mathcal{F}}R_{[\nu]}^{1/3} \left(1 + \frac{\pi^3}{96} \frac{C_{\mathcal{F}}^2}{R_{[\nu]}^{2/3}}\right)$ .

*Proof.* Write  $\nu_j \bar{\omega} = Re^{i\varphi_j}$ ,  $j = 1, 2, 3, 4$ , with  $\varphi_1 < \varphi_2 < \varphi_3 < \varphi_4 \leq \varphi_1 + \pi/2$  so that  $\text{Arc}[\nu] = (\varphi_4 - \varphi_1)R$ , and note that  $\varphi_1 + \varphi_2 + \varphi_3 + \varphi_4 = 0$  by the definition of  $\omega$ . Therefore

$$C_{\mathcal{F}([\nu])} = \frac{2R^{2/3} |\text{Im}((\nu_1 - \nu_4)\bar{\omega})|}{(2\text{Re}(\sigma\bar{\omega}))^{1/3}} = \frac{2R^{2/3} |\sin(\varphi_1) - \sin(\varphi_4)|}{|2(\cos(\varphi_1) + \cos(\varphi_2) + \cos(\varphi_3) + \cos(\varphi_4))|^{1/3}}.$$

Now  $\sin(\varphi_1) - \sin(\varphi_4) = 2 \sin(\frac{\varphi_1 - \varphi_4}{2}) \cos(\frac{\varphi_1 + \varphi_4}{2})$  and  $\cos(\varphi_1) + \cos(\varphi_2) + \cos(\varphi_3) + \cos(\varphi_4) = 2(\cos(\frac{\varphi_1 - \varphi_4}{2}) + \cos(\frac{\varphi_2 - \varphi_3}{2})) \cos(\frac{\varphi_1 + \varphi_4}{2})$  since  $\frac{\varphi_2 + \varphi_3}{2} = -\frac{\varphi_1 + \varphi_4}{2}$ , and therefore

$$(6.2) \quad C_{\mathcal{F}} = \frac{2R^{2/3} |\sin(\frac{\varphi_1 - \varphi_4}{2})| |\cos(\frac{\varphi_1 + \varphi_4}{2})|^{2/3}}{|\frac{1}{2}(\cos(\frac{\varphi_1 - \varphi_4}{2}) + \cos(\frac{\varphi_2 - \varphi_3}{2}))|^{1/3}} \leq \frac{2R^{2/3} |\sin(\frac{\varphi_1 - \varphi_4}{2})|}{|\cos(\frac{\varphi_1 - \varphi_4}{2})|^{1/3}}$$

since  $0 \leq \cos(\frac{\varphi_4 - \varphi_1}{2}) \leq \cos(\frac{\varphi_3 - \varphi_2}{2})$ , as  $0 \leq \varphi_3 - \varphi_2 \leq \varphi_4 - \varphi_1 \leq \pi/2$ . By Lemma 6.2 we deduce that  $C_{\mathcal{F}} < 1.0106(\varphi_4 - \varphi_1)R^{2/3} = 1.0106R^{-1/3}\text{Arc}[\nu]$  and part (i) follows.

If  $(\varphi_4 - \varphi_1) - (\varphi_3 - \varphi_2) \geq 0.137(\varphi_4 - \varphi_1)^3$ , then  $C_{\mathcal{F}} < (\varphi_4 - \varphi_1)R^{2/3} = R^{-1/3}\text{Arc}[\nu]$  by Lemma 6.2 (ii), and the result follows. Otherwise  $(\varphi_4 - \varphi_3) + (\varphi_2 - \varphi_1) < 0.137(\varphi_4 - \varphi_1)^3$ ; and suppose, for example, that  $\varphi_2 - \varphi_1 < \frac{0.137}{2}(\varphi_4 - \varphi_1)^3$ . Applying the argument in the proof of Theorem 1.2 to the triangle formed by  $\nu_1, \nu_2, \nu_4$  we obtain

$$2R \leq 4\Delta R = |\nu_1 - \nu_2||\nu_1 - \nu_4||\nu_2 - \nu_4| \leq \text{Arc}(\nu_1, \nu_2)\text{Arc}(\nu_1, \nu_4)\text{Arc}(\nu_2, \nu_4) = (\varphi_2 - \varphi_1)(\varphi_4 - \varphi_1)(\varphi_4 - \varphi_2)R^3 < 0.0685(\varphi_4 - \varphi_1)^5 R^3 = 0.0685R^{-2}\text{Arc}^5[\nu],$$

so that  $\text{Arc}[\nu] > 1.963667195R^{3/5} > C_{\mathcal{F}}R^{1/3}$  for  $R > 0.08C_{[\nu]}^{15/4}$  and part (ii) follows.

Let us suppose that  $\varphi_4 - \varphi_1 = 4\lambda$  so that  $\pi/2 \geq 4\lambda \geq 0$ . Therefore  $0 = \varphi_1 + \varphi_2 + \varphi_3 + \varphi_4 = 4\varphi_1 + m\lambda$  for some  $m$ ,  $4 \leq m \leq 12$ , so that  $\varphi_1 + \varphi_4 = 2\varphi_1 + 4\lambda = (4 - m/2)\lambda$  and therefore  $|(\varphi_1 + \varphi_4)/2| \leq \lambda$ . We therefore deduce that  $|\cos(\frac{\varphi_1 + \varphi_4}{2})| \geq |\cos \lambda| \geq \cos^2 \lambda = (1 + \cos 2\lambda)/2 \geq |\frac{1}{2}(\cos(\frac{\varphi_1 - \varphi_4}{2}) + \cos(\frac{\varphi_2 - \varphi_3}{2}))|$ . Therefore (6.2) implies that

$$(6.3) \quad C_{\mathcal{F}} \geq 2R^{2/3} \left| \sin\left(\frac{\varphi_1 - \varphi_4}{2}\right) \right| \left| \cos\left(\frac{\varphi_1 - \varphi_4}{2}\right) \right|^{1/3}.$$

Applying Lemma 6.2(iii) yields  $C_{\mathcal{F}} \geq R^{2/3} \frac{2}{\pi} |\varphi_1 - \varphi_4| = R^{-1/3} \frac{2}{\pi} \text{Arc}[\nu]$  so that

$$(6.4) \quad \text{Arc}[\nu] < (\pi/2)C_{\mathcal{F}}R^{1/3},$$

and also that

$$C_{\mathcal{F}} \geq 2R^{2/3} \left( \frac{\varphi_1 - \varphi_4}{2} - \frac{(\varphi_1 - \varphi_4)^3}{24} \right) = \frac{\text{Arc}[\nu]}{R^{1/3}} - \frac{\text{Arc}^3[\nu]}{12R^{7/3}} \geq \frac{\text{Arc}[\nu]}{R^{1/3}} - \frac{\pi^3}{96} \frac{C_{\mathcal{F}}^3}{R^{4/3}}$$

which implies part (iii).  $\square$

## 7. CLASSIFICATION

**Theorem 7.1.** *If  $\text{Arc}[\nu] < tR^{1/3}$  where  $[\nu]$  is primitive and non-degenerate, then either  $\text{Arc}[\nu] \geq \frac{\pi}{2}R_{[\nu]}$  with  $R \leq (2t/\pi)^{3/2}$ , or  $[\nu] \in \mathcal{F}$ , for some  $\mathcal{F} \in \mathcal{F}(1.01062t)$  with  $R \leq 0.084t^{15/4}$ , or  $[\nu] \in \mathcal{F}$  for some  $\mathcal{F} \in \mathcal{F}(t)$ .*

*Proof.* If  $\text{Arc}[\nu] \geq \frac{\pi}{2}R$  then  $\frac{\pi}{2}R < tR^{1/3}$  and the first option follows. If  $C_{\mathcal{F}} < t$  then the second option follows. Finally suppose that  $\text{Arc}[\nu] < \frac{\pi}{2}R_{[\nu]}$  and  $C_{\mathcal{F}} \geq t$ . Then  $tR^{1/3} > \text{Arc}[\nu] > 0.9895C_{[\nu]}R^{1/3}$  by Theorem 6.1 i), so that  $C_{\mathcal{F}} \leq 1.01062t$ , that is  $\mathcal{F}_{[\nu]} \in \mathcal{F}(1.01062t)$ . But then  $R \leq 0.084t^{15/4}$  else  $R > 0.084t^{15/4} > (0.08)(1.01062t)^{15/4} \geq 0.08C_{\mathcal{F}}^{15/4}$  implying that  $\text{Arc}[\nu] > C_{\mathcal{F}}R^{1/3} \geq tR^{1/3}$  by Theorem 6.1 ii).  $\square$



*Proof of Theorem 1.4.*  $\max\{2^{-18t^{15}}, 0.084t^{15/4}, (2t/\pi)^{3/2}\} = 2^{-18t^{15}}$  for  $t \geq 2.432362919\dots$ , and is  $< 2.4$  for smaller  $t$ . The result then follows from Theorem 7.1 and Corollary 4, and by verifying that the 4-tuples of lattice points on the circles of radius  $\leq \sqrt{5}$  satisfy our bound.  $\square$

**Lemma 7.1.** *Given  $[\nu] \in \mathcal{F}$  there exists  $[\nu]_n$  such that  $R_{[\nu]_n}^2 \leq C_{\mathcal{F}}^3 p_{[\nu]}^3$ .*

*Proof.* Fix  $\delta > 0$ , and select  $m$  such that  $\text{Arc}([\nu]_m) \leq (1 + \delta)C_{\mathcal{F}}R^{1/3}$ , which is possible by (3.9). For convenience we replace  $[\nu]$  by  $[\nu]_m$ . By (3.2) we have, using the arithmetic-geometric mean inequality,

$$\begin{aligned} R_{[\nu]_n}^2 / R_{[\nu]}^2 &= \prod_{i=1}^3 \left( \alpha^{2n} \left| \frac{\omega_i + \bar{\omega}_i}{2} \right|^2 + \beta^{2n} \left| \frac{\omega_i - \bar{\omega}_i}{2} \right|^2 \right) \\ &\leq \left( \frac{\alpha^{2n} \sum_{i=1}^3 \left| \frac{\omega_i + \bar{\omega}_i}{2} \right|^2 + \beta^{2n} \sum_{i=1}^3 \left| \frac{\omega_i - \bar{\omega}_i}{2} \right|^2}{3} \right)^3 \\ &\leq \left( \alpha^{2n} + \beta^{2n} (1 + \delta)^2 \frac{C_{\mathcal{F}}^2}{8R^{4/3}} \right)^3 \end{aligned}$$

To obtain this last inequality we first note that each  $\left| \frac{\omega_i + \bar{\omega}_i}{2} \right| \leq 1$  and that, if we write each  $\nu_j = Re^{i\varphi_j}$  where  $\varphi_1 \leq \varphi_2 \leq \varphi_3 \leq \varphi_4 < \varphi_1 + \pi/2$  then  $2|\omega_1 - \bar{\omega}_1| = 4|\sin(\frac{\varphi_1 - \varphi_2 - \varphi_3 + \varphi_4}{4})| \leq |\varphi_1 - \varphi_2 - \varphi_3 + \varphi_4| = |(\varphi_4 - \varphi_1) - (\varphi_3 - \varphi_1) - (\varphi_2 - \varphi_1)| \leq |\varphi_4 - \varphi_1|$  and, similarly,  $2|\omega_2 - \bar{\omega}_2| \leq |\varphi_4 - \varphi_1|$  and  $|\omega_3 - \bar{\omega}_3| \leq |\varphi_4 - \varphi_1|$ , so that

$$\frac{1}{3} \sum_{i=1}^3 \left| \frac{\omega_i - \bar{\omega}_i}{2} \right|^2 \leq \frac{(\varphi_4 - \varphi_1)^2}{8} = \frac{\text{Arc}[\nu]^2}{8R^2} \leq (1 + \delta)^2 \frac{C_{\mathcal{F}}^2}{8R^{4/3}}.$$

Now let  $n$  be the integer closest to  $\frac{\log((1+\delta)^2 C_{\mathcal{F}}^2 / (8R^{4/3}))}{4 \log \alpha}$ ; in fact suppose that this equals  $n - \gamma$  with  $|\gamma| \leq 1/2$ . Then

$$R_{[\nu]_n}^2 \leq R^2 \left( (1 + \delta) \frac{C_{\mathcal{F}}}{8^{1/2} R^{2/3}} (\alpha^{2\gamma} + |\beta|^{2\gamma}) \right)^3 \leq \frac{(1 + \delta)^3 C_{\mathcal{F}}^3 (\alpha + |\beta|)^3}{8^{3/2}}.$$

Now  $\alpha + |\beta| = 2p$  if  $\epsilon = 1$ , and  $\alpha + |\beta| = 2q\sqrt{Q}$  if  $\epsilon = -1$ . In the latter case we have  $q\sqrt{Q} = p + 1/(p + q\sqrt{Q}) \leq 2^{1/2}p$  (which is attained when  $Q = 2$ ). We obtain our result by an appropriate choice of  $\delta$ , since  $R_{[\nu]_n}^2$  is an integer.  $\square$

## 8. OUR ALGORITHM

Fix  $t > 0$  and suppose that we wish to determine all primitive 4-tuples  $[\nu]$  of lattice points for which  $\text{Arc}[\nu] \leq tR_{[\nu]}^{1/3}$ .

**Step 0. Small 4-tuples.** We examine all  $[\nu]$  such that  $R_{[\nu]} \leq (2t/\pi)^{3/2}$ .

**Step 1. Finding admissible  $Q$ .** We determine all positive integers  $Q \not\equiv 0 \pmod{4}$ , whose prime factors are 2 or are  $\equiv 1 \pmod{4}$ , which can be written as the product of four co-prime integers all of which are  $\leq t^3/16$  (by Lemmas 4.1 and 4.3).

**Step 2. Degenerate 4-tuples.** By Lemma 5.1 we examine, for each such square  $Q$ , all  $[\nu]$  with  $Q_{[\nu]} = Q$  and  $R_{[\nu]} \leq (t/2)^6 Q^{3/4}$ .

**Step 3. Finding the families.** For each such non-square  $Q$ , consider all the families containing a 4-tuples  $[\nu]$  with  $R_{[\nu]}^2 \equiv 0 \pmod{Q_1}$ , where  $Q_1 = Q/(2, Q)$ , such that  $R_{[\nu]}^2 \leq (1.01062t)^3 p_{[\nu]}^3$  (by Lemma 7.1 and theorem 6.1 (i)).

**Step 4. Computing the constants.** Compute the constants  $C_{\mathcal{F}_{[\nu]}}$  for the families obtained in step 3.

**Step 5. Families with  $t \leq C_{\mathcal{F}} < 1.01062t$ .** We examine all  $[\nu] \in \mathcal{F}$  such that  $R_{[\nu]} \leq 0.08t^{15/4}$ . (Theorem 6.1 (ii))

**Step 6. Families with  $C_{\mathcal{F}} < t$ .** We examine all  $[\nu] \in \mathcal{F}$  such that  $R_{[\nu]} \leq \left(\frac{C_{\mathcal{F}}^3 \pi^3}{t - C_{\mathcal{F}}}\right)^{3/2}$ . (Theorem 6.1 (iii))

For example, suppose that we wish to determine all primitive 4-tuples  $[\nu]$  of lattice points for which  $\text{Arc}[\nu] \leq 5R_{[\nu]}^{1/3}$ : To begin with we determine all such  $[\nu]$  for which  $R_{[\nu]}^2 \leq (10/\pi)^3 < 33$ ; these all happen to be degenerate examples (see the table below). In step 1 we see that  $5^3/16 < 8$  and so the possible values of the four (pairwise co-prime) factors of  $Q$  are 1, 2 and 5, so that  $Q = 1, 2, 5$  or 10. In step 2 we look for degenerate examples on circles of radius  $\leq (5/2)^6 Q^{3/4}$ ,  $Q = 1, 2, 5, 10$  according, finding

$R^2$	$[\nu]$	$\text{Arc}[\nu]R^{-1/3}$
5	$1 + 2i, 2 + i, 2 - i, 1 - 2i$	<b>3.7863...</b>
65	$7 + 4i, 8 + i, 8 - i, 7 - 4i$	<b>4.1746...</b>
5	$2 + i, 2 - i, 1 - 2i, -1 - 2i$	<b>4.2716...</b>
25	$5i, 3 + 4i, 4 + 3i, 5$	<b>4.5930...</b>
13	$2 + 3i, 3 + 2i, 3 - 2i, 2 - 3i$	<b>4.6217...</b>
125	$10 + 5i, 11 + 2i, 11 - 2i, 10 - 5i$	<b>4.6364...</b>
325	$17 + 6i, 18 + i, 18 - i, 17 - 6i$	<b>4.6655...</b>
85	$2 + 9i, 6 + 7i, 7 + 6i, 9 + 2i$	<b>4.9836...</b>
533	$22 + 7i, 23 + 2i, 23 - 2i, 22 - 7i$	<b>4.9953...</b>

as well as examples equivalent to these via multiplication by  $1, -1, i$  or  $-i$ , or via complex conjugation.

In step 3, noting that  $p_{[\nu]} = 1$  if  $Q_{[\nu]} = 1$  or 2,  $p_{[\nu]} \in \{2, 1/2\}$  if  $Q = 5$  and  $p_{[\nu]} = 3$  if  $Q = 10$ , we consider those  $[\nu]$  for which  $R_{[\nu]}^2$  is divisible by 1, 5 or 5 respectively, while being  $\leq (5.06)^3, (10.11)^3$  or  $(15.16)^3$ , respectively. Then in step 4 we found seven families (which we describe on the table overleaf), with constants

$$\begin{aligned} C_{\mathcal{F}_1} &= 2 \left(\frac{5}{3}(3 + \sqrt{10})\right)^{\frac{1}{3}} < C_{\mathcal{F}_2} = (20)^{\frac{1}{3}} \left(\frac{1+\sqrt{5}}{2}\right) < C_{\mathcal{F}_3} = 4 \left(\frac{1}{7}(5 + 4\sqrt{2})\right)^{\frac{1}{3}} \\ &< C_{\mathcal{F}_4} = \left(\frac{10}{3}(14 + 5\sqrt{10})\right)^{\frac{1}{3}} < C_{\mathcal{F}_5} = 3\sqrt{2} \left(\frac{2}{7}(5 + 3\sqrt{2})\right)^{\frac{1}{3}} < C_{\mathcal{F}_6} = 10^{\frac{1}{3}} \left(\frac{1+\sqrt{5}}{2}\right)^{\frac{5}{3}} \\ &< C_{\mathcal{F}_7} = 2(1 + \sqrt{2}) < 5 < C_{\mathcal{F}_8} = \frac{3 \cdot 2^{\frac{2}{3}}}{5^{\frac{1}{6}}} \left(\frac{1+\sqrt{5}}{2}\right)^{\frac{4}{3}}. \end{aligned}$$

Step 5 is vacuous and, for step 6, a simple calculation reveals that  $\text{Arc}[\nu] \leq 5R_{[\nu]}^{1/3}$  for all  $[\nu]_n \in \mathcal{F}_m : 1 \leq m \leq 7$ , except the initial 4-tuples  $[\nu]_0$  in  $\mathcal{F}_1, \mathcal{F}_2, \mathcal{F}_5$  and  $\mathcal{F}_7$ .

*Proof of Corollary 2.* By Theorem 6.1(i) we see that  $\text{Arc}[\nu] \geq C_{\mathcal{F}_1} R_{[\nu]}^{1/3}$  if  $[\nu] \in \mathcal{F}_m : 2 \leq m \leq 7$  since  $C_{\mathcal{F}_1} < 0.9895C_{\mathcal{F}_2}$ . By Theorem 2 we see that  $\text{Arc}[\nu] \geq C_{\mathcal{F}_1} R_{[\nu]}^{1/3}$  for  $[\nu] \in \mathcal{F}_1$  if  $R_{[\nu]} > 19.7897\dots$ , which is the case for all  $[\nu] \in \mathcal{F}_1$  except the initial 4-tuple, but in that case  $\text{Arc}[\nu] \approx 5.3720\dots R_{[\nu]}^{1/3}$ . We are therefore left only with a subset of the degenerate cases given above, namely the top three cases, which each have  $R \leq \sqrt{65}$ .  $\square$

	$\mathcal{F}$	$[\nu]_0$	$[\nu]_1$	$[\nu]_2$	$[\nu]_3$	$[\nu]_4$
$\mathcal{F}_1$ <b>4.347370624..</b> $\sqrt{Q}\omega^2 = 1 + 3i$ $\alpha = 3 + \sqrt{10}$	$\mathcal{F}_4$	(1,-2) (2,1) (1,2) (-1,2)	(190,155) (197,146) (202,139) (206,133)	(46561,33448) (46520,33505) (46489,33548) (46463,33584)	(10882804,7844567) (10883057,7844216) (10883248,7843951) (10883408,7843729)	(2546700757,1835553826) (2546699198,1835555989) (2546698021,1835557622) (2546697035,1835558990)
$\mathcal{F}_2$ <b>4.392019964..</b> $\sqrt{Q}\omega^2 = 2 + i$ $\alpha = (1 + \sqrt{5})/2$	$\mathcal{F}_2$	(1,-2) (2,-1) (2,1) (-1,2)	(3,4) (4,3) (5,0) (4,-3)	(18,-1) (18,1) (17,6) (15,10)	(70,25) (71,22) (73,14) (74,7)	(308,59) (307,64) (304,77) (301,88)
$\mathcal{F}_3$ <b>4.601544787..</b> $\sqrt{Q}\omega^2 = 1 + i$ $\alpha = 1 + \sqrt{2}$	$\mathcal{F}_5$	(8,-1) (8,1) (7,4) (4,7)	(91,48) (93,44) (96,37) (99,28)	(1342,531) (1338,541) (1331,558) (1322,579)	(18739,7822) (18749,7798) (18766,7757) (18787,7706)	(264028,109219) (264004,109277) (263963,109376) (263912,109499)
$\mathcal{F}_4$ <b>4.631841066..</b> $\sqrt{Q}\omega^2 = 1 + 3i$ $\alpha = 3 + \sqrt{10}$	$\mathcal{F}_1$	(37,16) (35,20) (29,28) (28,29)	(7516,5483) (7532,5461) (7568,5411) (7573,5404)	(1766281,1272658) (1766183,1272794) (1765961,1273102) (1765930,1273145)	(413271130,297871475) (413271734,297870637) (413273102,297868739) (413273293,297868474)	(96707495041,69702803308) (96707491319,69702808472) (96707482889,69702820168) (96707481712,69702821801)
$\mathcal{F}_5$ <b>4.65445600..</b> $\sqrt{Q}\omega^2 = 1 + i$ $\alpha = 1 + \sqrt{2}$	$\mathcal{F}_3$	(1,-2) (2,-1) (2,1) (-2,1)	(6,7) (7,6) (9,2) (9,-2)	(113,36) (112,39) (108,49) (104,57)	(1528,659) (1531,652) (1541,628) (1549,608)	(21653,8906) (21646,8923) (21622,8981) (21602,9029)
$\mathcal{F}_6$ <b>4.804476431..</b> $\sqrt{Q}\omega^2 = -1 + 2i$ $\alpha = 2 + \sqrt{5}$	$\mathcal{F}_8$	(16,13) (13,16) (8,19) (5,20)	(774,1307) (789,1298) (810,1285) (821,1278)	(60800,98145) (60737,98184) (60648,98239) (60601,98268)	(4613294,7465447) (4613561,7465282) (4613938,7465049) (4614137,7464926)	(350706224,567450437) (350705093,567451136) (350703496,567452123) (350702653,567452644)
$\mathcal{F}_7$ <b>4.828427124..</b> $\sqrt{Q}\omega^2 = -1 + i$ $\alpha = 1 + \sqrt{2}$	$\mathcal{F}_7$	(-2,1) (-1,2) (2,1) (2,-1)	(9,8) (8,9) (1,12) (-1,12)	(48,149) (51,148) (68,141) (72,139)	(869,2018) (862,2021) (821,2038) (811,2042)	(11762,28589) (11779,28582) (11878,28541) (11902,28531)

Here  $\mathcal{F}_8 = \mathcal{F}(1 + 2i, -2 - i, 2 + i, 1 - 2i)$ .

## 9. ASYMPTOTICS

Fix  $t$ . We wish to determine the number of 4-tuples of lattice points which lie on an arc of length  $tR^{1/3}$  of a circle of radius  $R$  centered at the origin, where  $R \leq x$ . First we deal only with primitive 4-tuples. Since there only a bounded number of degenerate  $[\nu]$  with  $\text{Arc}[\nu] < tR^{1/3}$  by Corollary 4, this reduces to determining the number of primitive 4-tuples in each family  $\mathcal{F}$  with  $C_{\mathcal{F}} < t$ . For a given family  $\mathcal{F}$  we have  $R_n \sim \alpha^{3n} |\text{Re}(\sigma\bar{\omega})|/4$  by (3.5) so the number with  $R_n \leq x$  is  $\log x/3 \log \alpha + O(1)$ . Now each such 4-tuple is one of an equivalence class of 8 examples (as we have discussed). Therefore there are a total of  $\sim \beta_t \log x$  primitive 4-tuples of lattice points which lie on an arc of length  $tR^{1/3}$  on a circle of radius  $R \leq x$  centered at the origin, where

$$\beta_t = \frac{8}{3} \sum_{\mathcal{F}: C_{\mathcal{F}} < t} \frac{1}{\log \alpha(\mathcal{F})} = \frac{8}{3} \sum_{\alpha} \frac{|\mathcal{F}_{\alpha}(t)|}{\log \alpha},$$

with  $\mathcal{F}_{\alpha}(t) = \{\mathcal{F}, \alpha(\mathcal{F}) = \alpha, C_{\mathcal{F}} < t\}$ . Rather like in the prime number theorem, if we count each primitive 4-tuple  $[\nu]$  with weight  $\log \alpha = \log \alpha_{[\nu]}$  then we have

$$\sum_{\substack{\text{primitive } [\nu] \\ \text{Arc}[\nu] < tR^{1/3} \\ R_{[\nu]} < x}} \log \alpha_{[\nu]} \sim \frac{8}{3} \#\{\mathcal{F} : C_{\mathcal{F}} < t\} \log x.$$

Write

$$N_4(t, x) = \#\{[\nu], \text{Arc}[\nu] < tR_{[\nu]}^{1/3}, R_{[\nu]} \leq x\}$$

and, for  $g \in \mathbb{Z}[i]$ ,

$$N_4(t, x, g) = \#\{[\nu] = (\nu_1, \nu_2, \nu_3, \nu_4), \gcd(\nu_1, \nu_2, \nu_3, \nu_4) = g, \text{Arc}[\nu] < tR^{1/3}, R_{[\nu]} \leq x\}.$$

We proved above that  $N_4(t, x, 1) \sim \beta_t \log x$  for fixed  $t$  as  $x \rightarrow \infty$ . To estimate  $N(t, x)$  we use the formula  $\text{Arc}[\nu] = |g|\text{Arc}[\nu/g]$  to obtain

$$N_4(t, x) = \sum_g N_4(t, x, g) = \sum_g N_4(t|g|^{-2/3}, x|g|^{-1}, 1) \sim \sum_g \beta_{t|g|^{-2/3}} \log(x|g|^{-1}) \sim B_t \log x,$$

as  $x \rightarrow \infty$ , where

$$B_t = \frac{8}{3} \sum_{\alpha} \frac{1}{\log \alpha} \sum_g |\mathcal{F}_{\alpha}(t|g|^{-2/3})|,$$

and the sums are over all  $g = a + bi$ ,  $0 \leq b < a$  for which  $|g| \leq (t/C_{\mathcal{F}_1})^{3/2}$ .

By Theorem 1.4 and the fact that  $\mathcal{F}(t)$  is finite, we know that  $B_t$  is a piecewise constant function. We conjecture that  $B_t \asymp t^3 \log^6 t$  and, more generally, that

$$N_4(t, x) \asymp \min\{x^2, t^3\} (\log(\min\{x^2, t^3\}))^6 \log x$$

for all  $t > (40 + \frac{40}{3}\sqrt{10})^{1/3}$ . One can prove the slightly stronger estimate  $N_4(t, x) \sim c_4 x^2 \log^7 x$  for  $x \leq (t/(2\pi))^{3/2}$  for some constant  $c_4 > 0$  by a simple counting argument (the analogous argument for  $N_3$  is given at the beginning of the next section).

**9.1. Asymptotic 3-tuples.** Writing  $N_3(t, x) = \#\{3\text{-tuples } [\nu], |\nu| \leq x, \text{Arc}[\nu] < t|\nu|^{1/3}\}$ , we conjecture that

$$(9.1) \quad N_3(t, x) \asymp x^{2/3} \min\{x^2, t^3\}^{2/3} (\log(\min\{x^2, t^3\}))^3$$

for all  $t > 16^{1/3}$ . We can prove a slightly stronger result when  $x \leq (t/(2\pi))^{3/2}$ , since in this range all the 3-tuples with  $|\nu| \leq x$  are counted in  $N_3(t, x)$ , so that

$$N_3(t, x) = \sum_{n \leq x^2} \binom{r(n)}{3} = \frac{1}{6} \sum_{n \leq x^2} r^3(n) + O(r^2(n)) \sim c_3 x^2 \log^3 x,$$

for some constant  $c_3 > 0$ , via the usual counting argument using contour integration. The conjecture in (9.1) is equivalent to  $N_3(t, x) \asymp x^{2/3} t^2 \log^3 t$  for  $x \geq (t/(2\pi))^{3/2}$ ; we now prove a weak version of this estimate.

**Theorem 9.1.**

$$x^{2/3} t^2 \log t \ll N_3(t, x) \ll x^{2/3} t^2 \log^3 t, \quad x \gg t^{3/2} \gg 1.$$

*Proof.* Defining

$$N_3(t, x, g) = \#\{[\nu] = (\nu_1, \nu_2, \nu_3), \gcd(\nu_1, \nu_2, \nu_3) = g, |\nu| \leq x, \text{Arc}[\nu] < t|\nu|^{1/3}\}$$

we have

$$N_3(t, x) = \sum_{|g| \leq t^{3/2}} N_3(t, x, g) = \sum_{|g| \leq t^{3/2}} N_3(t|g|^{-2/3}, x|g|^{-1}, 1),$$

so to prove the theorem it suffices to obtain upper and lower bounds for primitive 3-tuples (that is, the case  $g = 1$ ), which we will do in lemmas 9.4 and 9.5 below.

First we introduce some lemmas and notation. For any  $l \in \mathbb{Z}[i]$  let  $\varphi_l$  be the argument of  $l$ , and then  $\mathbb{Z}_0[i] = \{l \in \mathbb{Z}[i], 0 \leq \varphi_l \leq \pi/2\}$  with  $\mathbb{Z}_0^v[i]$  the set of visible lattice points in  $\mathbb{Z}_0[i]$  (that is the numbers  $a + bi \in \mathbb{Z}[i]$  with  $(a, b) = 1$  and  $a \geq b \geq 0$ ).

**Lemma 9.1.** *For any 3-tuple  $[\nu]$  there exists unique  $g \in \mathbb{Z}_0[i]$ ,  $\mathbf{p}_1, \mathbf{p}_2, \mathbf{p}_3 \in \mathbb{Z}_0^v[i]$ ,  $(\mathbf{p}_i, \mathbf{p}_j) = 1$ , and  $t_1, t_2, t_3 \in \{0, 1, 2, 3\}$  such that*

$$\begin{aligned}\nu_1 &= i^{t_1} g \mathbf{p}_1 \mathbf{p}_2 \overline{\mathbf{p}_3} \\ \nu_2 &= i^{t_2} g \mathbf{p}_1 \overline{\mathbf{p}_2} \mathbf{p}_3 \\ \nu_3 &= i^{t_3} g \overline{\mathbf{p}_1} \mathbf{p}_2 \mathbf{p}_3.\end{aligned}$$

*Proof.* The values  $g = (\nu_1, \nu_2, \nu_3)$ ,  $\mathbf{p}_1 = (\nu_1/g, \nu_2/g, \overline{\nu_3/g})$ ,  $\mathbf{p}_2 = (\nu_1/g, \overline{\nu_2/g}, \nu_3/g)$ ,  $\mathbf{p}_3 = (\overline{\nu_1/g}, \nu_2/g, \nu_3/g)$  are the only one satisfying the conditions.  $\square$

**Lemma 9.2.** *If  $\text{Arc}[\nu] < s|\nu|^{1/3}$  and  $t_1 = t_2 = t_3$ , then  $|\varphi_{\mathbf{p}_i} - \varphi_{\mathbf{p}_j}| < (s/2)|\nu|^{-2/3}$  for  $i \neq j$ .*

*Proof.* Notice that  $\varphi_{\mathbf{p}_2} - \varphi_{\mathbf{p}_3} = \frac{1}{2}(\varphi_{\nu_1} - \varphi_{\nu_2}) \leq \frac{s}{2}|\nu|^{-2/3}$ , and the same argument works for the other differences.  $\square$

**Lemma 9.3.** *There are at most  $4w^2\epsilon + 1$  visible lattice points in the angular sector  $|z| \leq w$ ,  $|\varphi_z - \alpha| \leq \epsilon$ .*

*Proof.* Let  $P_1, \dots, P_n$  denote the visible lattice points in the sector ordered according to increasing argument. Each of the triangles  $(O, P_i, P_{i+1})$ ,  $i = 1, \dots, n-1$  are inside the angular sector, each has area  $\geq 1/2$ , and they are disjoint so that

$$(n-1) \cdot \frac{1}{2} \leq \text{Area of the angular sector} = 2\epsilon w^2$$

$\square$

**Lemma 9.4.** *If  $y > s^{3/2}$  then  $N_3(s, y, 1) \ll y^{2/3} s^2 \log^2 s$ .*

*Proof.* We will count 3-tuples  $[\nu]$  with  $|\nu| \leq y$ ,  $\text{Arc}[\nu] < s|\nu|^{1/3}$  and  $g = \gcd(\nu_1, \nu_2, \nu_3) = 1$ , and restrict our attention, in the notation of lemma 9.1, to the case where  $t_1 = t_2 = t_3$  (the other cases following by analogous arguments), and  $|\mathbf{p}_1| \leq |\mathbf{p}_2| \leq |\mathbf{p}_3|$  (the other cases following by re-arrangement of the  $\nu_i$ ). If  $2^{j_i} \leq |\mathbf{p}_i| < 2^{j_i+1}$  for  $i = 1, 2, 3$ , then  $j_1 \leq j_2 \leq j_3$ , and the condition  $|\nu| \leq y$  implies that  $j := j_1 + j_2 + j_3 \leq \log_2 y$ . Now

$$2^{j_3} \leq |\mathbf{p}_3| \leq |\nu_2 - \nu_3| \leq \text{Arc}[\nu] \leq s|\nu|^{1/3} = s|\mathbf{p}_1 \mathbf{p}_2 \mathbf{p}_3|^{1/3} \leq (2s)2^{j/3}$$

so that  $j_3 \leq \log_2(2s) + j/3$ ; and, similarly,  $j_1, j_2 \leq \log_2(2s) + j/3$ . Together these imply that

$$(9.2) \quad -2\log_2(2s) \leq j_i - j/3 \leq \log_2(2s) \text{ for } i = 1, 2, 3.$$

The condition  $\text{Arc}[\nu] < s|\nu|^{1/3}$  implies that  $|\varphi_{\mathbf{p}_i} - \varphi_{\mathbf{p}_1}| < s2^{-1-2j/3}$  for  $i = 2, 3$  by lemma 9.2), and therefore  $p_i$  is a visible lattice point in the angular sector  $|z| \leq 2^{j_i+1}$ ,  $|\varphi_{\mathbf{p}_i} - \varphi_{\mathbf{p}_1}| < s2^{-2j/3}$ . There are  $\ll 2^{2j_i} \cdot s2^{-2j/3}$  such lattice points by lemma 9.3, since  $1 \leq 2^{2j_2} 2^{-j} 2^{j_3} \leq 2^{2j_2} 2^{-j} (2s) 2^{j/3} \leq (2s) 2^{2j_2} 2^{-2j/3} \leq (2s) 2^{2j_3} 2^{-2j/3}$ . There are  $\ll 2^{2j_1}$  lattice points  $\pi$  with  $|\mathbf{p}_i| < 2^{j_i+1}$ , and hence the total number of such triples  $\mathbf{p}_1, \mathbf{p}_2, \mathbf{p}_3$  is  $\ll s^2 2^{2j/3}$ .

Now, for a given  $j$ , there are  $\ll (\log s)^2$  triples of integers  $j_1, j_2, j_3$  by (9.2), and so our count of lattice points is

$$\ll \sum_{j \leq \log_2 y} s^2 2^{2j/3} (\log s)^2 \ll y^{2/3} s^2 (\log s)^2,$$

as required.  $\square$

We now prove bounds in the other direction:

**Lemma 9.5.** *We have  $N_3(s, y, 1) \gg y^{2/3} s^2$  for  $s$  sufficiently large.*

*Proof.* We construct examples in lemma 9.1 with  $g = 1$  and each  $t_i = 0$ , so we want  $\mathbf{p}_1, \mathbf{p}_2, \mathbf{p}_3 \in \mathbb{Z}_0^v[i]$  with each  $|\mathbf{p}_i| \leq y^{1/3}$ , as well as  $(\mathbf{p}_i, \mathbf{p}_j) = 1$  and  $|\varphi_{\mathbf{p}_i} - \varphi_{\mathbf{p}_j}| < (s/2)y^{-2/3}$  for  $i \neq j$ . Consider the lattice points  $\mathbf{p} = a + bi \in \mathbb{Z}_0^v[i]$  with  $a + b$  odd,  $|\mathbf{p}| \leq y^{1/3}$ , and divide the circle of radius  $y^{1/3}$  into angular sectors of width  $(s/2)y^{-2/3}$ , which we will denote  $S_1, \dots, S_k$  with  $k \asymp [4\pi y^{2/3}/s]$ . Note that  $(\mathbf{p}, \bar{\mathbf{p}}) = 1$  as  $\mathbf{p}$  is visible and  $a + b$  is odd. There are  $\sim (1/2\pi)y^{2/3}$  such lattice points, and so an average of  $\sim s/8\pi^2$  per sector: we can thus prove that if  $s \geq 25\pi^2$  then there are  $\gg ks^3 \gg y^{2/3}s^2$  triples  $\mathbf{p}_1, \mathbf{p}_2, \mathbf{p}_3 \in \mathbb{Z}_0^v[i]$  with each  $|\mathbf{p}_i| \leq y^{1/3}$ , and  $|\varphi_{\mathbf{p}_i} - \varphi_{\mathbf{p}_j}| < (s/2)y^{-2/3}$  for  $i \neq j$ ; however we do not necessarily have  $(\mathbf{p}_i, \mathbf{p}_j) = 1$  for each  $i \neq j$ . Forcing this to happen, complicates our argument.

We modify the above construction by only considering those  $\mathbf{p} = a + bi \in \mathbb{Z}_0^v[i]$  with  $a + b$  odd,  $|\mathbf{p}| \leq y^{1/3}$ , which have no divisor  $g \in \mathbb{Z}[i]$  for which  $1 < |g| \leq B$ , for some large fixed  $B$  to be chosen later. Sieve methods yield that there are  $\asymp y^{2/3}/\log B$  such lattice points. Thus if  $(\mathbf{p}_i, \mathbf{p}_j) = g \neq 1$  then  $|g| > B$ . Now this implies that  $\mathbf{p}_i/g, \mathbf{p}_j/g$  are distinct visible lattice points lying in an angular sector with  $|z| \leq y^{1/3}/|g|$  of angular width  $(s/2)y^{-2/3}$ : There are  $\leq 1 + 2(s/2)y^{-2/3}(y^{1/3}/|g|)^2 = 1 + s/|g|^2$  such points by lemma 9.3, and thus no such pair if  $|g|^2 > s$ , and at most  $k(1 + s/|g|^2)(s/|g|^2)/2 \leq ks^2/|g|^4$  such pairs otherwise. Therefore the total number of pairs of such lattice points  $\mathbf{p}_i, \mathbf{p}_j$  with  $|(\mathbf{p}_i, \mathbf{p}_j)| > B$  is

$$\leq \sum_{g: B < |g| \leq \sqrt{|s|}} \frac{ks^2}{|g|^4} \leq \frac{8ks^2}{B^2} \leq \frac{101y^{2/3}s}{B^2}.$$

To complete our proof we translate this into a graph theory problem: Let  $G_i$  denote a graph whose vertices are the  $\mathbf{p} = a + bi \in \mathbb{Z}_0^v[i]$  with  $a + b$  odd,  $|\mathbf{p}| \leq y^{1/3}$ , which have no divisor  $g \in \mathbb{Z}[i]$  for which  $1 < |g| \leq B$ , and where  $\mathbf{p} \in S_i$ . Two vertices in  $G_i$  have an edge between them if the corresponding  $\mathbf{p}$  have no common factor. Then  $N_3(s, y, 1)$  is at least the total number of triangles in all of the graphs  $G_i$ . Let  $n_i$  be the number of vertices in  $G_i$ , and let  $e_i$  be the number of edges in the complement of  $G_i$ ; that is the number of pairs of  $\mathbf{p}$  in this sector that have a common factor. Then

$$n := n_1 + \dots + n_k \asymp y^{2/3}/\log B, \quad k \asymp y^{2/3}/s, \quad e := e_1 + \dots + e_k \ll y^{2/3}s/B^2.$$

Our result follows from the following by taking  $B$  and then  $s$  sufficiently large.

**Lemma 9.6.** *Fix  $\epsilon > 0$ . There exist constants  $c, C > 0$  such that for any integers  $n, e, k$  satisfying  $n \geq Ck$  and  $e \leq cn^2/k$ , if  $G_i$  is a graph created by deleting  $e_i$  edges from the complete graph on  $n_i \geq 3$  vertices, for  $i = 1, 2, \dots, k$ , then there are  $\geq (1/4 - \epsilon)n^3/k^2$  triangles in the set of graphs  $G_1, \dots, G_k$ , where  $n := n_1 + \dots + n_k$  and  $e := e_1 + \dots + e_k$ .*

*Proof.* A graph  $G_i$  with  $n_i^2/4 + m_i$  edges contains  $\gg m_i n_i$  distinct triangles. Hence the total number of triangles in  $G_1, \dots, G_k$  is

$$(9.3) \quad \gg \sum_{i=1}^k n_i \max \left\{ 0, \frac{n_i^2 - 2n_i}{4} - e_i \right\}.$$

We shall suppose that for given  $n, e, k$  we have the choice of non-negative real numbers  $e_i$  and  $n_1 \geq n_2 \geq \dots \geq n_k \geq 3$  that minimizes the right side of (9.3). This evidently minimized by taking  $e_i = e_{i,0} := \frac{n_i(n_i-2)}{4}$  for  $i = 1, 2, \dots, \ell - 1$ , with  $0 \leq e_\ell \leq e_{\ell,0}$ , and  $e_i = 0$  for  $i > \ell$ . Now, for fixed  $e - e_\ell = \sum_{1 \leq i < \ell} \frac{n_i(n_i-2)}{4}$ , we wish to maximize  $\sum_{1 \leq i < \ell} n_i$  (so as to minimize  $\sum_{i > \ell} n_i$ ), and thus we take them all to be equal. If  $n_\ell > n_k$  with  $e_\ell < e_{\ell,0}$  then we get a contradiction of minimality by taking  $n'_1 = n_1 - \delta$ ,  $n'_k = n_k + \delta$  for some very small  $\delta > 0$ .

Thus we have  $n_1 = \dots = n_{\ell-1} = r$ , say, and  $n_\ell = \dots = n_k = s$ , say, with  $r \geq s \geq 3$ . Moreover we have  $0 < e - (\ell - 1)\frac{r(r-2)}{4} \leq \frac{s(s-2)}{4}$  and  $r(\ell - 1) + s(k - \ell + 1) = n$ , and that the right side of (9.3) equals  $(k + 1 - \ell)\frac{s^2(s-2)}{4} - e_\ell s$ .

This is now a classical optimization problem. If we take  $\ell - 1 = \lambda k$  for  $0 \leq \lambda \leq 1$  and  $r = \rho n/k, s = \sigma n/k$  then  $\rho\lambda + \sigma(1 - \lambda) = 1$ , so that  $\rho \geq 1 \geq \sigma \geq 0$ . Solving we find that  $\sigma = (1 - \rho\lambda)/(1 - \lambda)$ . Now  $e = \frac{r(r+O(1))}{4}(\ell + O(1)) = \lambda\rho^2 n^2/4k\{1 + O(1/r + 1/\ell)\}$  so that  $\lambda \leq 4c/\rho^2\{1 + O(1/r + 1/\ell)\}$ , and so  $\sigma > 1 - O(c)$ . The quantity to be minimized is  $\geq (1 - \lambda)ks^3/4\{1 + O(1/s)\} = (1 - O(c))\sigma^3 n^3/4k^2 \geq (1 - O(c))n^3/4k^2$ . implying the desired result.  $\square$

**Remark 2.** *By slightly modifying this proof one can show that if  $n \geq (2 + \epsilon)k$  and  $e < (1/4 - \epsilon)n(n - 2k)/k$  then there are  $\gg_\epsilon n^3/k^2$  triangles. Evidently there can be no triangles if  $n \leq 2k$  (if every  $G_i$  has degree at most two), nor if  $e > n^2/4k$  (by making each  $G_i$  a complete bipartite graph with about  $n/2k$  vertices in each class).*

## 10. CLOSE DIVISORS

In a forthcoming paper [5], we will deal with the analogous problem for close divisors  $d_1, \dots, d_k$  of a rational integer  $N$ . The quantity  $\omega = (d_1 \cdots d_k)^{1/k}/N^{1/2}$  shows where the lattice points  $(d_i, N/d_i)$  are located on the hyperbola  $xy = N$ . Since each  $N/d_i$  is also a divisor of  $N$ , we study only the  $k$ -tuples of *large* divisors, that is those with  $\omega \geq 1$ . We can give a lower bound for  $L(d_1, \dots, d_k) = \max_{i \neq j} |d_i - d_j|$ , analogous to theorem 1.1.

$$(10.1) \quad L(d_1, \dots, d_k) \geq N^{1/4 - 1/(8[(k-1)/2] + 4)}.$$

The analysis of the cases  $k = 2, 3$  is similarly straightforward and one can obtain the sharp estimates:

$$L(d_1, d_2) \geq 2, \quad L(d_1, d_2, d_3) \geq 2^{2/3}N^{1/6};$$

and that  $\{L(d_1, d_2, d_3)N^{-1/6}\}$  is dense in  $[2^{2/3}, \infty)$ .

The case  $k = 4$  again requires more delicate arguments, as we found in this article for the analogous problem for lattice points on circles. For the problem of close divisors, (10.1) yields  $L(d_1, \dots, d_4) \geq N^{1/6}$ , and the exponent “1/6” cannot be increased as we see from the following example: The integers  $N_n = 2p_n p_{n+1} p_{n+2} q_n q_{n+1} q_{n+2}$ , have divisors

$$d_{1,n} = 2q_n p_{n+1} q_{n+2}, \quad d_{2,n} = 2q_n q_{n+1} p_{n+2}, \quad d_{3,n} = p_n p_{n+1} p_{n+2}, \quad d_{4,n} = 2p_n q_{n+1} q_{n+2},$$

where  $(p_n, q_n)$  denote the solutions of the Pell equation  $x^2 - 2y^2 = \pm 1$ . In fact

$$\lim_{n \rightarrow \infty} L(d_{1,n}, d_{2,n}, d_{3,n}, d_{4,n})N_n^{-1/6} = 2^{7/12} + 2^{13/12} = 3.6172\dots$$

This is not the only family with this kind of property. As in this paper, we can classify all the “close” 4-tuples  $[d] = (d_1, d_2, d_3, d_4)$  of divisors of integers  $N$  into families  $\mathcal{F}$  such that

$$\lim L(d_{1,n}, d_{2,n}, d_{3,n}, d_{4,n})N_n^{-1/6} = C_{\mathcal{F}},$$

One can determine a formula similar to (3.7) to describe each family. We are then able to deduce the following theorem (analogous to Corollary 2 herein):

**Theorem 10.1.** *For any large 4-tuple of divisors  $d_1, d_2, d_3, d_4$  of  $N$  we have*

$$L(d_1, d_2, d_3, d_4) > 2 \left(\frac{3}{2}\right)^{1/12} \left(\frac{8 + 3\sqrt{6}}{5}\right)^{1/3} N^{1/6},$$

whenever  $N > N_0$ . On the contrary there exist infinitely many integers large 4-tuple of divisors  $d_{1,n}, d_{2,n}, d_{3,n}, d_{4,n}$  of  $N_n$  with

$$\lim_{n \rightarrow \infty} \frac{L(d_{1,n}, d_{2,n}, d_{3,n}, d_{4,n})}{N_n^{1/6}} = 2 \left( \frac{3}{2} \right)^{1/12} \left( \frac{8 + 3\sqrt{6}}{5} \right)^{1/3} = 3.006555939..$$

Although the ideas and techniques used in [5] are similar to those used in this paper, there are sufficient differences that it seems necessary to write a different paper.

We do not know if the exponent in (10.1) is sharp for  $k \geq 5$ , just as we do not know if the exponent in Theorem 1.1 is sharp for  $k \geq 5$ .

## 11. OTHER RELATED QUESTIONS

Herein we have studied very precise questions on close lattice points on a circle; and in [5] we develop a similar study of close lattice points on a hyperbola. Presumably it should be possible to generalize these results to close lattice points on all other curves of degree two in the plane, and perhaps to curves of higher degree. In this case one knows that there are very few points, after Mumford's theorem and Faltings' theorem, and those that there are should presumably be very sparse, but such questions appear, for now, to lie deep. There is an important school of research that attempts to obtain bounds which are within a small factor of best possible, which makes these bounds very applicable. Like in proofs of Theorem 1.1, the key articles by Bombieri and Pila [1], Heath-Brown [9], and then Elkies [7], all use combinatorial arguments and linear algebra; these have the severe limitation that they are unlikely to give bounds for typical curves that are much better than what is obtained for the lattice-point rich, rational curve,  $y = x^d$ . Quite recently, Ellenberg and Venkatesh [8] have incorporated true arithmetic-geometric techniques into these arguments, so as to distinguish between rational and non-rational curves, and thus they get bounds of a strength that had previously seemed inaccessible.

One can also ask about analogous questions in higher dimension, for instance how close can one pack  $k$  lattice points on a sphere in  $\mathbb{R}^3$ ? One has to be a little careful as Heath-Brown showed us: Select integer  $r$  which has many representations as the sum of two squares; for example, if  $r$  is the product of  $\ell$  distinct primes that are  $\equiv 1 \pmod{4}$  then  $r$  has  $2^\ell$  such representations. Now let  $N$  be an arbitrarily large integer and consider the set of representations of  $n = N^2 + r$  as the sum of three squares. Evidently we have  $\geq 2^\ell$  such representations in an interval whose size, which depends only on  $\ell$ , is independent of  $n$ . Note though that these lattice points all lie on the hyperplane  $x = N$ , so we can better formulate our question by asking: *How close can one pack  $k$  lattice points on a sphere in  $\mathbb{R}^3$ , no four of which belong to the same hyperplane?*

## 12. OPEN PROBLEMS

We finish this article with two open problems.

**Problem 1:** *Do there exist infinitely many circles  $x^2 + y^2 = R_n^2$  with five lattice points on an arc of length  $\ll R_n^{2/5}$ ?*

We doubt it: From Theorem 1.1 we know that an arc of length  $R^{2/5}$  contains, at most, four lattice points, and our guess is that the exponent  $2/5$  can be increased, perhaps to as much as  $1/2$ .



**Problem 2:** *Is there a uniform bound for the number of lattice points on an arc of length  $\ll R^{1/2}$ ?*

Theorem 1.1 gives the upper bound  $\ll \log R$ , for the number of lattice points on an arc of length  $R^{1/2}$ , and we would like to see this significantly improved. Indeed this provokes the following (see also [4]):

**Conjecture 1.** *For every  $\epsilon > 0$  there exists a constant  $B_\epsilon$ , such that there are no more than  $B_\epsilon$  lattice points on an arc of length  $R^{1-\epsilon}$  of a circle of radius  $R$  that is centered at the origin.*

Fix  $m$ . Let  $a$  be a large integer. Let  $\{\sigma_\ell : j = 1, 2, \dots, \binom{2m}{m}\}$  be the set of functions  $\sigma_\ell : \{1, \dots, 2m\} \rightarrow \{-1, 1\}$  for which  $\sum_{j=1}^{2m} \sigma_\ell(j) = 0$ . Now define  $v_\ell := \prod_{j=1}^{2m} (a + j + i\sigma_\ell(j))$ . Obviously  $v_\ell = \prod_{j=1}^{2m} (a + j)(1 + O(m^2/a^2))$ , so that  $|v_i - v_j| \ll_m |v_i|^{1-1/m}$ ; in other words we have constructed  $\binom{2m}{m}$  lattice points in an arc of length  $O(r^{1-1/m})$  on a circle of radius  $r$ . Thus if Conjecture 1 is true then  $B_\epsilon$  would have to be at least  $e^{C/\epsilon}$  for some constant  $C > 0$ .

#### REFERENCES

- [1] Enrico Bombieri and Jonathan Pila, The number of integral points on arcs and ovals. *Duke Math. J.* **59** (1989), 337–357.
- [2] Javier Cilleruelo, Arcs containing no three lattice points. *Acta Arithmetica* **59** (1991), 87–90.
- [3] Javier Cilleruelo and Antonio Córdoba, Trigonometric polynomials and lattice points. *Proc. Amer. Math. Soc.* **115**, (1992), 899–905.
- [4] Javier Cilleruelo and Andrew Granville, Lattice points on circle, Squares in arithmetic progressions and Sumset of squares *Proc. Workshop “Additive Combinatorics”, Montreal (2006)*.
- [5] Javier Cilleruelo and Andrew Granville, Close divisors. *In preparation*.
- [6] Javier Cilleruelo and Gerald Tenenbaum, An overlapping theorem with applications. *Publicacions Matemàtiques* to appear.
- [7] Noam D. Elkies, Rational points near curves and small nonzero  $|x^3 - y^2|$  via lattice reduction, in Algorithmic Number Theory (Leiden, 2000), *Lecture Notes in Comput. Sci.* **1838** (2000), 33–63, Springer-Verlag, New York.
- [8] Jordan Ellenberg and Akshay Venkatesh, On uniform bounds for rational points on nonrational curves. *Int. Math. Res. Not.* **35** (2005), 2163–2181.
- [9] Roger Heath-Brown The density of rational points on curves and surfaces. *Ann. of Math.* **155** (2002), 553–595.
- [10] Antony Zygmund, A Cantor-Lebesgue theorem for double trigonometric series. *Studia Mathematica* **43** (1972), 173–178.

DEPARTAMENTO DE MATEMÁTICAS. UNIVERSIDAD AUTÓNOMA DE MADRID. 28049 MADRID  
*E-mail address:* franciscojavier.cilleruelo@uam.es

DÉPARTEMENT DE MATHÉMATIQUES ET STATISTIQUE, UNIVERSITÉ DE MONTRÉAL, CP 6128 SUCC  
 CENTRE-VILLE, MONTRÉAL, QC H3C 3J7, CANADA  
*E-mail address:* andrew@dms.umontreal.ca