

Bookmarks Menu

[Two new fungus gnats \(Insecta, Diptera, Mycetophilidae\) from the Lower Cretaceous of Spain](#)
[The vertebrate assemblage of Buenache de la Sierra \(Upper Barremian of Serranía de Cuenca, Spain\) with insights into its taphonomy and palaeoecology. Histological observations of Enantiornithine bone \(Saurischia, Aves\) from the Lower Cretaceous of Las Hoyas \(Spain\)](#)
[Trace fossils and their palaeoecological significance in Lower Cretaceous lacustrine conservation deposits, El Montsec, Spain](#)
[Early Cretaceous Ferns From Lacustrine Limestones At Las Hoyas, Cuenca Province, Spain](#)
[At the feet of the dinosaurs: the early history and radiation of lizards](#)
[An unusual lizard \(Reptilia: Squamata\) from the Early Cretaceous of Las Hoyas, Spain](#)
[The fish trace fossil *Undichna* from the Cretaceous of Spain](#)
[A new species of *Mirovia* \(Coniferales, Miroviaceae\) from the Lower Cretaceous of the Iberian Ranges \(Spain\)](#)
[Frenelopsis \(Coniferales: Cheirolepidiaceae\) And Related Male Organ Genera From The Lower Cretaceous Of Spain](#)
[Conclusiones / Bibliografía](#)
[Phylogenetic relationships of fossil and Recent gonorynchiform fishes \(Teleostei: Ostariophys\)](#)
[Molecular taphonomy of macrofossils from the Cretaceous Las Hoyas Formation, Spain](#)
[A phylogenetic system of Clavatoraceae \(fossil Charophyta\)](#)
[Taphonomie des plantes et interprétations paléocéologiques. Une synthèse](#)
[Taphonomy of insects in carbonates and amber](#)
[On the aerodynamics of leading-edge high-lift devices of avian wings](#)
[Mesozoic chrysopid-like Planipennia: a phylogenetic approach \(Insecta: Neuroptera\)](#)
[A new assemblage of Spanish Early Cretaceous teleostean fishes, formerly considered “leptolepids”: phylogenetic relevance](#)
[First isotopic and multidisciplinary evidence for nonmarine coelacanths and pycnodontiform fishes: palaeoenvironmental implications](#)
[New Cretaceous Scoliidae \(Vespida=Hymenoptera\) from the Lower Cretaceous of Spain and Brazil](#)
[A Nestling Bird from the Lower Cretaceous of Spain: Implications for Avian Skull and Neck Evolution](#)
[COPTOCLAVID BEETLES \(COLEOPTERA: ADEPHAGA\) FROM THE LOWER CRETACEOUS OF SPAIN: A NEW FEEDING STRATEGY IN BEETLES](#)
[Xeromorphic adaptations of some Mesozoic gymnosperms. A review with palaeoclimatological implications](#)
[A New Crested Pterosaur from the Early Cretaceous of Spain: The First European Tapejarid \(Pterodactyloidea: Azhdarchoidea\)](#)
[The Birds from Las Hoyas](#)
[Phylogeny and classification of the Stenophlebioptera \(Odonata: Epiproctophora\)](#)
[Anurans from the Early Cretaceous Lagerstätte of Las Hoyas, Spain: New evidence on the Mesozoic diversification of crown-clade Anura](#)
[Palaeohistology and preservation of tetrapods from Las Hoyas \(Lower Cretaceous, Spain\)](#)
[Leaf architecture and ecophysiology of an early basal eudicot from the Early Cretaceous of Spain: Early eudicot leaf architecture and ecophysiology. Scrutinizing Barremian coprolite inclusions to record digestive strategies](#)
[Exceptional coprolite association from the Early Cretaceous continental Lagerstätte of Las Hoyas, Cuenca, Spain](#)
[A tiny lizard \(Lepidosauria, Squamata\) from the Lower Cretaceous of Spain: TINY FOSSIL LIZARD ARTICLE](#)
[The Gobiosuchidae in the early evolution of Crocodyliformes](#)
[Did all theropods have chicken-like feet? First evidence of a non-avian dinosaur podotheca](#)
[A presumed spelaeogriphacean crustacean from an upper Barremian wetland \(Las Hoyas; Lower Cretaceous; Central Spain\): SPELAEOGRIPHACEAN FROM LAS HOYAS](#)
[Las Hoyas: a Cretaceous wetland: a multidisciplinary synthesis after 25 years of research on an exceptional fossil Lagerstätte from Spain](#)
[Montsechia, an ancient aquatic angiosperm](#)
[INVOLVEMENT OF MICROBIAL MATS IN DELAYED DECAY: AN EXPERIMENTAL ESSAY ON FISH PRESERVATION](#)
[Involvement of microbial mats in early fossilization by decay delay and formation of impressions and replicas of vertebrates and invertebrates](#)
[The Effect Of microbial Mats In The Decay Of Anurans With Implications For Understanding Taphonomic Processes In The Fossil Record](#)
[A diminutive perinate European Enantiornithes reveals an asynchronous ossification pattern in early birds](#)
[An articulated hindlimb of a basal iguanodont \(Dinosauria, Ornithomoda\) from the Early Cretaceous Las Hoyas Lagerstätte \(Spain\)](#)
[Feather diversity in the Barremian \(Early Cretaceous\) of Las Hoyas, Spain](#)
[Soft-tissue and dermal arrangement in the wing of an Early Cretaceous bird: Implications for the evolution of avian flight](#)
[A bizarre, humped Carcharodontosauria \(Theropoda\) from the Lower Cretaceous of Spain](#)
[First fossil harvestmen \(Arachnida: Opiliones\) from Spain and notes on the fossil record of Opiliones](#)
[An isolated bird foot from the Barremian \(Lower Cretaceous\) of Las Hoyas \(Cuenca, Spain\)](#)
[An Early Cretaceous faunal and floral continental assemblage:Las Hoyas fossil site \(Cuenca, Spain\)](#)
[Soft-bodied annelids \(Oligochaeta\) from the Lower Cretaceous \(La Huérguina Formation\) of the Las Hoyas Konservat-Lagerstätte, Spain](#)
[First pterosaur remains from the Early Cretaceous Lagerstätte of Las Hoyas, Spain: palaeoecological significance](#)
[Dental characters and phylogeny of pycnodontiform fishes](#)
[Polazzodus, gen. nov., a new pycnodont fish from the Late Cretaceous of northeastern Italy](#)
[Sylvienodus, a new replacement genus for the Cretaceous pycnodontiform fish “Pycnodus” laveirensis](#)
[A Cretaceous eutriconodont and integument evolution in early mammals](#)
[Anatomical differentiation of isolated scales of amiiform fishes \(Amiiformes, Actinopterygii\) from the Early Cretaceous of Las Hoyas \(Cuenca, Spain\)](#)
[A new genus and species for the amiiform fishes previously assigned to Amiopsis from the Early Cretaceous of Las Hoyas, Cuenca, Spain](#)
[The Exceptional Fossil Site of Las Hoyas \(SPAIN\) from an Educational Perspective](#)
[A new trackway possibly made by a trotting theropod at the Las Hoyas fossil site \(Early Cretaceous, Cuenca Province, Spain\): Identification, bio-dynamics and palaeoenvironmental implications](#)
[Fully fledged enantiornithine hatchling revealed by Laser-Stimulated Fluorescence supports precocial nesting behavior](#)
[Reassignment of Lioestheria codoensis Cardoso \(Spinicaudata, Anthronestheriidae\) from the Lower Cretaceous of Brazil: systematics and paleoecology.](#)
[Palaeobiology of tanaidaceans \(Crustacea: Peracarida\) from Cretaceous ambers: extending the scarce fossil record of a diverse peracarid group](#)
[The Stratigraphy and Rifting Evolution of the Oxfordian–Barremian \(Upper Jurassic–Lower Cretaceous\) in the Serranía de Cuenca \(Southwestern Iberian Ranges, Spain\)](#)
[The Karstic Habitat of Spelaeogriphaceans from the Las Hoyas Fossil Site \(Upper Barremian, Serranía de Cuenca, Spain\)](#)
[Palaeoenvironmental Interpretation of Palaeosols and Palustrine Carbonates of the Earliest Cretaceous Terrestrial Ecosystems in the Serranía de Cuenca, Iberian Ranges, Spain](#)
[Short-term colonization trace-fossil assemblages in a carbonate lacustrine Konservat-Lagerstätte \(Las Hoyas fossil site, Lower Cretaceous, Cuenca, Central Spain\)](#)
[The enigmatic Mesozoic insect taxon Chresmodidae \(Polyneoptera\): New palaeobiological and phylogenetic data, with the description of a new species from the Lower Cretaceous of Brazil](#)
[New Leptotarsus from the Early Cretaceous of Brazil and Spain: the oldest members of the family Tipulidae \(Diptera\)](#)
[The Larvae of the Mesozoic family Aeschniidae and their Phylogenetic Implications \(Insecta, Odonata, Anisoptera\)](#)
[A comparative overview of the neuropteran assemblage of the Lower Cretaceous Yixian Formation \(China\), with description of a new genus of Psychopsidae \(Insecta: Neuroptera\)](#)
[A new species of Tipulidae \(Diptera\) from the Lower Cretaceous Yixian Formation of Liaoning, China – Evolutionary implications](#)
[New Unionida \(Bivalvia\) from the Kimmeridgian \(Late Jurassic\) of Asturias, Spain, and their palaeobiogeographical implications](#)
[Presence of organic layers in shells of fossil and recent Unionida \(Bivalvia\) and their implications](#)
[Late Jurassic biogeochemical microenvironments associated with microbialite-coated unionids \(Bivalvia\), Asturias \(N Spain\)](#)
[Floristic and vegetational changes in the Iberian Peninsula during Jurassic and Cretaceous](#)
[Effects of chemical preparation protocols on δ13C values of plant fossil samples](#)
[Reassessment of the cheirolepidiacean conifer Frenelopsis teixeirae Alvin et Pais from the Early Cretaceous \(Hauterivian\) of Portugal and palaeoenvironmental considerations](#)
[Woody or not woody? Evidence for early angiosperm habit from the Early Cretaceous fossil wood record of Europe](#)
[Early Angiosperm Ecology: Evidence from the Albian-Cenomanian of Europe](#)
[Vertebrate diversity of the Jehol Biota as compared with other lagerstätten](#)
[Mineralogía y cambios composicionales en fragmentos óseos atribuidos a un dinosaurio ornitópodo del yacimiento barremiense de Buenache de la Sierra \(Formación Calizas de La Huérguina, Cuenca, España\)](#)
[THE IMPACT OF MICROBIAL MATS AND THEIR MICROENVIRONMENTAL CONDITIONS IN EARLY DECAY OF FISH](#)
[Preservation in microbial mats: mineralization by a talc-like phase of a fish embedded in a microbial sarcophagus](#)
[Palaeoclimate, environmental factors, and bird body size: A multivariable analysis of avian fossil preservation](#)
[The mineralization of dinosaur soft tissue in the Lower Cretaceous of Las Hoyas, Spain](#)
[A New Freshwater Basal Eucryptodiran Turtle from the Early Cretaceous of Spain](#)
[A sinemydid turtle from the Jehol Biota provides insights into the basal divergence of crown turtles](#)
[A metamorphosed salamander from the early Cretaceous of Las Hoyas, Spain](#)
[Phylogeny And Systematics Of Squamata \(Reptilia\) Based On Morphology](#)
[A new Early Cretaceous salamander \(Regalerpeton weichangensis gen. et sp. nov.\) from the Huajjying Formation of northeastern China](#)
[Exploring Patterns and Extent of Bias in Estimating Divergence Time from Mitochondrial DNA Sequence Data in a Particular Lineage: A Case Study of Salamanders \(Order Caudata\)](#)
[Notas nomenclaturales sobre anfibios actuales y fósiles](#)
[Fossils, molecules, divergence times, and the origin of Salamandroidea](#)
[Late Jurassic salamandroid from western Liaoning, China](#)
[A new albanerpetontid amphibian from the Early Cretaceous \(Barremian\) Wessex Formation of the Isle of Wight, southern England](#)
[Xenopus in Space and Time: Fossils, Node Calibrations, Tip-Dating, and Paleobiogeography](#)
[Jeholotriton paradoxus \(Amphibia: Caudata\) from the Lower Cretaceous of southeastern Inner Mongolia, China](#)
[A Late Jurassic salamander \(Amphibia: Caudata\) from the Morrison Formation of North America](#)
[Ontogeny Discombobulates Phylogeny: Paedomorphosis and Higher-Level Salamander Relationships](#)
[Largest bird from the Early Cretaceous and its implications for the earliest avian ecological diversification](#)
[The origin and early evolution of birds: discoveries, disputes, and perspectives from fossil evidence](#)
[A new, three-dimensionally-preserved enantiornithine bird \(Aves: Ornithothoraces\) from Gansu Province, north-western China: NEW ENANTIORNITHIAN FROM GANSU](#)
[The phylogenetic position of Ambiortus: Comparison with other Mesozoic birds from Asia](#)
[Second species of enantiornithine bird from the Lower Cretaceous Changma Basin, northwestern China with implications for the taxonomic diversity of the Changma avifauna](#)
[A historical specimen of enantiornithine bird from the Early Cretaceous of Mongolia representing a new taxon with a specialized neck morphology](#)
[The first dentary of Lisboasaurus \(Crocodylomorpha, ?Mesoeucrocodylia\) from the Lower Cretaceous \(Barremian\) of Uña, Cuenca Province, Spain](#)
[Particulate emission from biomass combustion](#)
[A new semionotiform actinopterygian fish from the Mesozoic of Spain and its phylogenetic implications](#)
[A Berriasian actinopterygian fauna from Cherves-de-Cognac, France: Biodiversity and palaeoenvironmental implications](#)
[Cenomanian transgression in the Basque-Cantabrian Basin \(northern Spain\) and associated faunal replacement](#)
[Akromystax tilmachiton gen. et sp. nov., a new pycnodontid fish from the Lebanese Late Cretaceous of Haql and En Nammoura](#)
[Local-scale analysis of plant community from the Early Cretaceous riparian ecosystem of Hautrage, Belgium](#)
[δ13C and stomatal number variability in the Cretaceous conifer Frenelopsis](#)
[\(2301\) Proposal to conserve the name Frenelopsis against Aularthrophyton \(fossil Gymnospermae: Coniferales\)](#)
[A holistic approach to the palaeoecology of Las Hoyas Konservat-Lagerstätte \(La Huérguina Formation, Lower Cretaceous, Iberian Ranges, Spain\)](#)
[The Posture of Tyrannosaurus rex : Why Do Student Views Lag Behind the Science?](#)
[New Eoptychoptera \(Insecta: Diptera, Ptychopteridae\) from the Lower Cretaceous of Spain](#)
[Pectoral girdle and forelimb variation in extant Crocodylia: the coracoid-humerus pair as an evolutionary module: Variation of the Crocodylian Skeleton](#)
[Geometric morphometrics in macroevolution: morphological diversity of the skull in modern avian forms in contrast to some theropod dinosaurs](#)
[Disparity and geometry of the skull in Archosauria \(Reptilia: Diapsida\): ARCHOSAURIAN SKULL DISPARITY](#)
[Constraining uncertainty in the timescale of angiosperm evolution and the veracity of a Cretaceous Terrestrial Revolution](#)
[Canrightia resinifera gen. et sp. nov., a new extinct angiosperm with Retimonocolpites -type pollen from the Early Cretaceous of Portugal: missing link in the eumagnoliid tree?](#)
[Barremian Angiosperm Pollen and Associated Palynomorphs from the Dakhla Oasis Area, Egypt](#)
[The ancestral flower of angiosperms and its early diversification](#)
[Fossil evidence for Cretaceous escalation in angiosperm leaf vein evolution](#)
[Deciphering Early Angiosperm Landscape Ecology Using a Clustering Method on Cretaceous Plant Assemblages](#)
[Influence of latitude and climate on spread, radiation and rise to dominance of early angiosperms during the Cretaceous in the Northern Hemisphere](#)
[Rise to dominance of angiosperm pioneers in European Cretaceous environments](#)
[CO2 and temperature decoupling at the million-year scale during the Cretaceous Greenhouse](#)
[Taphonomy and palaeoecology in the upper Barremian of the SW Iberian chain \(Spain\): A model to compare taxonomy and diversity of biotas from different coeval basins](#)
[Plant Tissue Decay in Long-Term Experiments with Microbial Mats](#)
[Inferring flight parameters of Mesozoic avians through multivariate analyses of forelimb elements in their living relatives](#)
[Flight reconstruction of two European enantiornithines \(Aves, Pygostylia\) and the achievement of bounding flight in Early Cretaceous birds](#)
[No evidence for directional evolution of body mass in herbivorous theropod dinosaurs](#)
[The first definitive Asian spinosaurid \(Dinosauria: Theropoda\) from the early cretaceous of Laos](#)
[European ornithomimosaur \(Dinosauria, Theropoda\): an undetected record](#)
[Cranial Osteology of Haplocheirus sollers Choiniere et al., 2010 \(Theropoda: Alvarezsauroidea\)](#)
[A New Troodontid Theropod from the Late Cretaceous of Central China, and the Radiation of Asian Troodontids](#)
[THE DIET OF OSTRICH DINOSAURS \(THEROPODA: ORNITHOMIMOSAURIA\)](#)
[A new North American therizinosaurid and the role of herbivory in ‘predatory’ dinosaur evolution](#)
[A giant ornithomimosaur from the Early Cretaceous of China](#)
[New information on Nqwebasaurus thwazi, a coelurosaurian theropod from the Early Cretaceous Kirkwood Formation in South Africa](#)
[The cranial osteology of Concavenator corcovatus \(Theropoda: Carcharodontosauria\) from the Lower Cretaceous of Spain](#)
[6. Mesozoic mammals — early mammalian diversity and ecomorphological adaptations](#)
[From the Field to the Museum: Analysis of Groups-Purposes-Locations in Relation to Spain’s Moveable Palaeontological Heritage](#)
[The decline of farmland birds in Spain is strongly associated to the loss of fallowland](#)
[A novel approach for the metric analysis of fern fronds: Growth and architecture of the Mesozoic fern Weichselia reticulata in the light of modern ferns](#)
[Palaeobiogeographic and metric analysis of the Mesozoic fern Weichselia](#)
[Helminth eggs from early cretaceous faeces](#)